

Centre for Ecology & Hydrology

Air Pollution: Deposition to and impacts on vegetation in (South-)East Europe, Caucasus, Central Asia (EECCA/SEE) and South-East Asia

Editors: Harry Harmens and Gina Mills ICP Vegetation Programme Coordination Centre, March 2014

D



Working Group on Effects of the Convention on Long-range Transboundary Air Pollution



Air Pollution: Deposition to and impacts on vegetation in (South-)East Europe, Caucasus, Central Asia (EECCA/SEE) and South-East Asia

Report prepared by the ICP Vegetation¹ March, 2014

Harry Harmens and Gina Mills (Editors)

ICP Vegetation Programme Coordination Centre,

Centre for Ecology and Hydrology,

Environment Centre Wales,

Deiniol Road, Bangor, Gwynedd, LL57 2UW, UK

Tel: + 44 (0) 1248 374500, Fax: + 44 (0) 1248 362133, Email: hh@ceh.ac.uk

http://icpvegetation.ceh.ac.uk

¹ International Cooperative Programme on Effects of Air Pollution on Natural Vegetation and Crops.

Acknowledgements

We wish to thank the UK Department for Environment, Food and Rural Affairs (Defra) for the continued financial support of the ICP Vegetation (Contract AQ0816). The United Nations Economic Commission for Europe (UNECE) and the UK Natural Environment Research Council (NERC) are also thanked for contributions to the funding of the ICP Vegetation Programme Coordination Centre. We wish to thank all the contributors to this report (listed below), ICP Vegetation participants that submitted country reports (see Annex 1) and ICP Vegetation participants who have provided data in the past that have been used in this report.

Harry Harmens	Centre for Ecology and Hydrology,	United Kingdom
Gina Mills	Bangor	
Katrina Sharps		
Lisa Emberson	Stockholm Environment Institute –	United Kingdom
Patrick Büker	York, University of York	
Joanne Morris		
Max Posch	Coordination Centre for Effects,	The Netherlands
Jaap Slootweg	RIVM, Bilthoven	
Jean-Paul Hettelingh		
Ilia Ilyin	EMEP/MSC-East	Russian Federation
Michael Gauss	EMEP/MSC-West	Norway
Anna Benedictow		

Contributors to this report

This report should be quoted as: Harmens, H. and Mills, G. (Eds.) (2014). Air Pollution: Deposition to and impacts on vegetation in (South-)East Europe, Caucasus, Central Asia (EECCA/SEE) and South-East Asia. ICP Vegetation Programme Coordination Centre, Centre for Ecology and Hydrology, Bangor, UK.

Cover photos were supplied by Patrick Büker, Martin Forsius, Felicity Hayes, Ilia Ilyin (map), Katrina Sharps (map), Norbert Schnyder, Dimitris Velissariou and Shutterstock (UK),

This report is the Copyright of Defra and has been prepared by CEH under contract to Defra (AQ0816). The contents of this report may not be reproduced in whole or in part, nor passed to any organisation or person without the specific prior written permission of Defra. CEH accepts no liability whatsoever to any third party for any loss or damage arising from any interpretation or use of the information contained in this report, or reliance on any views expressed therein. The NERC and CEH trademarks and logos ('the Trademarks') are registered trademarks of NERC in the UK and other countries, and may not be used without the prior written consent of the Trademark owner.

Executive summary

Increased ratification of the Protocols of the Convention on Long-range Transboundary Air Pollution (LRTAP) was identified as a high priority in the new long-term strategy of the Convention. Increased ratification and full implementation of air pollution abatement policies is particularly desirable for countries of Eastern Europe, the Caucasus and Central Asia (EECCA) and South-Eastern Europe (SEE). Hence, scientific activities within the Convention will need to involve these countries. In the current report, the ICP Vegetation has reviewed current knowledge on the deposition of air pollutants to and their impacts on vegetation in EECCA (Armenia, Azerbaijan, Belarus, Georgia, Kazakhstan, Kyrgyzstan, Moldova, Russian Federation, Tajikistan, Turkmenistan, Ukraine and Uzbekistan) and SEE countries (Albania, Bosnia and Herzegovina, Bulgaria, Croatia, Cyprus, Greece, Macedonia, Montenegro, Romania, Serbia, Slovenia and Turkey). As an outreach activity to Asia we have also reviewed current knowledge on this subject for the Malé Declaration countries in South-East Asia (SEA; Bangladesh, Bhutan, India, Iran, Maldives, Nepal, Pakistan and Sri Lanka). Air pollution is a main concern in Asia due to enhanced industrialisation, which is directly linked to continued strong economic growth in recent decades.

In these regions, there is generally a lack of an extensive network of monitoring stations to assess the magnitude of air concentrations and depositions of pollutants. In addition, emission inventories are often incomplete or not reported at all for some pollutants, which makes it difficult to validate atmospheric transport models for these regions. Furthermore, there is often a lack of coordinated monitoring networks to assess the impacts of air pollution on vegetation. Hence, the risk of adverse impacts on vegetation often has to be assessed using atmospheric transport models in conjunction with metrics developed to compute the risk of air pollution impacts on vegetation, such as critical loads and levels. Here we have focussed on the following air pollutants: nitrogen, ozone, heavy metals, POPs (EECCA/SEE countries) and aerosols, including black carbon as a component (South-East Asia).

Nitrogen

Critical load exceedances for nutrient nitrogen are only available for a limited number of EECCA countries. Compared with Western and Central Europe, available computed critical load exceedances for nitrogen have historically been lower in SEE and large areas of the EECCA region, particularly the northern part, and this was also the case in 2010. However, the critical load is expected to still be exceeded in large areas in 2020 with improvements since 2005 generally being lower than in Western and Central Europe, particularly in the EECCA region. Nitrogen concentrations in mosses were found to be intermediate to high in SEE compared to other European countries, indicating potentially a higher risk of nitrogen effects on ecosystems than computed by the critical loads. Little data is available on nitrogen deposition and impacts on vegetation in South-East Asia.

Ozone

For the first time, the ICP Vegetation has mapped the risk of adverse impacts of ozone on vegetation for the extended EMEP domain using the flux-based metric POD_Y^2 . The concentration-based approach (AOT40³) identifies the southern part of the EECCA region at highest risk, whereas the biologically more relevant flux-based approach (POD_Y) identifies the south-western part of the region bordering with Central Europe at highest risk. Both approaches indicate that the northern part of the region is at lowest risk of adverse impacts from ozone pollution. Field-based evidence is available for ozone impacts on crops in Greece and Slovenia, with many crop species showing visible leaf injury in Greece. Both AOT40 and POD_Y are computed to be high in SEE, indicating that this area is at high risk of ozone damage to vegetation. Staple food crops (maize, rice, soybean and wheat) are sensitive to moderately sensitive to ozone, threatening global food security. Recent flux-based risk assessment of ozone-induced wheat yield loss show that the estimated relative yield loss was 6.4-14.9% for China and 8.2-22.3% for India in 2000, with higher yield losses predicted for 2020, indicating the urgent

² Phytotoxic ozone dose above a flux threshold of Y nmol m⁻² projected leaf area s⁻¹

³ The accumulated hourly mean ozone concentration above 40 ppb, during daylight hours

need for curbing the rapid increase in surface ozone concentrations in this region. Worryingly, yield reductions of 20-35% have been recorded for various crop species when comparing yield in clean air with that in current ambient ozone concentrations in South-East Asia.

Heavy metals

Generally, deposition of heavy metals has declined in recent decades in EECCA and SEE countries, in agreement with the general decline computed for the rest of Europe, with the highest decline being reported for lead. However, apart from the western part of the EECCA region, the decline has generally been lower for EECCA and SEE countries compared to the rest of Europe. In 2011, the highest levels of metal deposition were computed in SEE countries, the south-western part and some (south-)eastern parts of the EECCA region. This might explain the relatively high concentrations of many heavy metals in mosses in countries in SEE Europe compared to the rest of Europe in recent years. High critical load exceedances have been reported for Macedonia for cadmium and lead and for Bosnia-Herzegovina and Russian Federation for lead. Widespread exceedance of the critical load for mercury has been observed in this region, similar to the rest of Europe. In India, the deposition of many heavy metals onto fruit and vegetables has been found to exceed WHO and Indian national limits for safe consumption.

Persistent organic pollutants (POPs)

Model assessment indicate a reduction of POP pollution in most of the EECCA and SEE countries between 1990 to 2011, particularly for hexachlorobenzene (HCB), although generally lower than for the rest of Europe; highest reductions were observed in the western part of the region. In 2011, the highest deposition for benzo[a]pyrene and polychlorinated dibenzodioxin were computed in the southwestern part of the region, whereas HCB levels were high for large parts of the Russian Federation.

Aerosols

South Asia is a region with high aerosol load compared to other regions, due its rapid growth and the arid climate. In particular the Indo-Gangetic Plain, South Asia's most important agricultural region, persistently has very high aerosol load, reducing visibility as well as solar radiation reaching the surface. Reduced photosynthesis might occur as a result of reduced solar radiation and larger aerosols blocking leaf pores, although the increase in diffuse radiation might have the opposite effect.

Conclusions and recommendations

This review highlights the lack of monitoring data regarding the deposition to and impacts of air pollutants on vegetation in EECCA/SEE countries and South-East Asia. It would be desirable to further enlarge coordinated networks to measure air concentrations and depositions of air pollutants, i.e. to extend the EMEP monitoring network in the EECCA/SEE region and establish a similar network in South-East Asia, for example by extending the Acid Deposition Monitoring Network in East Asia (EANET) by including other regions and more pollutants. International Cooperative Programmes of the LRTAP Convention might consider further stimulating the development of coordinated networks in these regions with the aim to establish widespread monitoring networks assessing the impacts of air pollutants on ecosystems. More measurement data are urgently needed to validate model outputs regarding the concentrations, deposition and associated risk for impacts of air pollutants on vegetation. The successful implementation of air pollution abatement policies in many other parts of Europe has highlighted the slower progress made with some of the air pollution abatement in the Eastern Europe, the Caucasus, Central and South-East Asia. Improvement of air quality in these regions will also benefit the rest of Europe due to a reduction on long-range transport of air pollutants, particularly those of hemispheric nature such as ozone and mercury. Many air pollution issues are remaining in the studied areas that require urgent attention, especially in regions of fast economic and population growth, ensuring future sustainable development without significant impacts on the functionality of ecosystems, the services they provide and food production.

Contents

Executive summary	3
1. Introduction	7
1.1 Background	7
1.2 ICP Vegetation	7
1.3 Aim and structure of the report	8
2. Air pollution deposition and impacts on vegetation in EECCA and SEE	9
2.1 Introduction	9
2.2 Nitrogen	10
2.2.1 Background	10
2.2.2 Nitrogen critical load exceedances	11
2.2.3 Nitrogen concentrations in mosses	12
2.3 Ozone	13
2.3.1 Background	13
2.3.2 Field-based evidence of ozone impacts	14
2.3.3 Modelled ozone risk assessment	15
2.4 Heavy metals	
2.4.1 Background	
2.4.2 Modelled heavy metal deposition	17
2.4.3 Concentrations of heavy metals in mosses	
2.4.4 Critical load exceedances for heavy metals	21
2.5 Persistent organic pollutants (POPs)	23
2.5.1 Background	23
2.5.2 Modelled deposition of POPs	23
2.5.3 POPs concentrations in mosses	24
3. Concentrations and effects of air pollutants on vegetation in	
South-East Asia	25
3.1 Background	25
3.2 Emission sources	25
3.3 Atmospheric concentration and deposition	
3.3.1 Ozone	
3.3.2 Nitrogen	
3.3.3 Atmospheric aerosols	30
3.3.4 Heavy metals	

3.4 Effects on vegetation
3.4.1 Ozone impacts
3.4.2 Heavy metal impacts
3.4.3 Nitrogen and sulphur impacts
3.4.4 Aerosol impacts
4. Conclusions and recommendations45
4.1 Conclusions
4.1.1 Eastern Europe, Caucasus and Central Asia (EECCA)
4.1.2 South-East Europe (SEE)45
4.1.3 South-East Asia (SEA; Malé Declaration countries)
4.2 Recommendations 46
References
References
References 47 Annex 1: Country reports 51 Albania 52
References 47 Annex 1: Country reports 51 Albania 52 Croatia 54
References 47 Annex 1: Country reports 51 Albania 52 Croatia 54 Egypt* 56
References 47 Annex 1: Country reports 51 Albania 52 Croatia 54 Egypt* 56 Greece 59
References 47 Annex 1: Country reports 51 Albania 52 Croatia 54 Egypt* 56 Greece 59 Macedonia 61
References 47 Annex 1: Country reports 51 Albania 52 Croatia 54 Egypt* 56 Greece 59 Macedonia 61 Romania 63
References 47 Annex 1: Country reports 51 Albania 52 Croatia 54 Egypt* 56 Greece 59 Macedonia 61 Romania 63 Russian Federation 65
References 47 Annex 1: Country reports 51 Albania 52 Croatia 54 Egypt* 56 Greece 59 Macedonia 61 Romania 63 Russian Federation 65 Serbia 68

* As outreach to North Africa

1. Introduction

Harry Harmens, Gina Mills

1.1 Background

Since being established in 1979, the Convention on Long-range Transboundary Air Pollution (LRTAP) has delivered demonstrable improvements in air quality resulting in, for example, reduced acidification of the environment, reductions in the highest peak levels of ozone and photochemical smog, and reductions in atmospheric heavy metal concentration and deposition. The LRTAP Convention has also begun to make improvements in atmospheric levels and deposition of nitrogen. Despite such good progress, air pollution in the UNECE region still causes significant environmental and health problems and new problems are emerging. Major strenghts and successes of the Convention are:

- Science-based policy decision making and the effects-oriented approach;
- Its geographical coverage of most of the northern hemisphere;
- The multi-pollutant/multi-effects approach of the Gothenburg Protocol on acidification, eutrophication (primarily nitrogen enrichment) and ground-level ozone, established in 1999;
- Establishment of the Protocols on Persistent Organic Pollutants (POPs) and on Heavy Metals in 1998, leading the way for a wider global approach to these problems, i.e. development of the Stockholm Convention on POPs and Minamata Convention on Mercury.

In 2010, the Convention adopted its long-term strategy for the next 10 years (EB Decision 2010/18). The strategy identified remaining challenges facing the UNECE region and emphasised that the focus of the work of the Convention should be on ozone, nitrogen and particulate matter, the latter primarily in relation to adverse impacts on human health. The Convention will pursue initiatives in addressing the synergies and trade-offs between policies to address air pollution, climate change and biodiversity. For example, there is a growing interest in the so-called short-lived climate forcers (SLCFs) as a potential means of mitigating short-term climate change before the effects of the longer-lived greenhouse gases are seen. SLCFs such as black carbon (a component of particulate matter) and ozone are being addressed in the revision of the Gothenburg Protocol, adopted in 2012.

Increased ratification of the Protocol on Heavy Metals, the Protocol on POPs and the Gothenburg Protocol was identified as a high priority in the new long-term strategy of the Convention. A viable future for the Convention depends on positive and vigorous participation by the Parties in all parts of the region, and on ensuring its extensive geographical coverage. Increased ratification and full implementation of air pollution abatement policies is particularly desirable for countries of Eastern Europe, the Caucasus and Central Asia (EECCA) and South-Eastern Europe (SEE). Hence, scientific activities within the Convention will need to involve these countries. In the current report, the ICP Vegetation has reviewed the knowledge on the deposition of air pollutants and their impacts on vegetation in EECCA (Armenia, Azerbaijan, Belarus, Georgia, Kazakhstan, Kyrgyzstan, Moldova, Russian Federation, Tajikistan, Turkmenistan, Ukraine and Uzbekistan) and SEE countries (Albania, Bosnia and Herzegovina, Bulgaria, Croatia, Cyprus, Greece, Macedonia, Montenegro, Romania, Serbia, Slovenia and Turkey). As an outreach activity to Asia we have also reviewed the current knowledge on this subject for the Malé Declaration countries in South-East Asia (Bangladesh, Bhutan, India, Iran, Maldives, Nepal, Pakistan and Sri Lanka).

1.2 ICP Vegetation

The International Cooperative Programme on Effects of Air Pollution on Natural Vegetation and Crops (ICP Vegetation) was established in 1987, initially with the aim to assess the impacts of air pollutants on crops, but in later years also on (semi-)natural vegetation. The ICP Vegetation is led by the UK and has its Programme Coordination Centre at the Centre for Ecology and Hydrology (CEH) in Bangor. The ICP Vegetation is one of seven ICPs and Task Forces that report to the Working Group on Effects (WGE) of the Convention on Long-range Transboundary Air Pollution (LRTAP Convention) on the effects of atmospheric pollutants on different components of the environment (e.g. forests,

fresh waters, materials) and health in Europe and North-America. The ICP Vegetation comprises an enthusiastic group of scientists from more than 40 countries (Harmens et al., 2013b), including scientists from outside the UNECE region as the ICP Vegetation stimulates outreach activities to other regions in the world. **Table 1.1** provides an overview of EECCA, SEE and Asian countries currently participating in the ICP Vegetation.

EECCA	SEE	Asia
Belarus (HM)	Albania (HM)	China (O ₃)
Russian Federation (HM)	Bulgaria (HM, N)	India (HM, O ₃)
Ukraine (HM, O ₃)	Croatia (HM, N, O ₃)	Japan (O ₃)
	Greece (O ₃)	Pakistan (O ₃)
	Macedonia (HM, N)	
	Romania (HM)	
	Serbia (HM)	
	Slovenia (HM, N, O ₃ , POPs)	
	Turkey (HM, N)	

Table 1.1EECCA, SEE^a and Asian countries participating in the ICP Vegetation. Within brackets:
pollutant(s) studied in the country.

^a Kosovo (United Nations administered territory, Security Council resolution 1244 (1999)) also participates (HM). HM = moss survey on heavy metals; N = moss survey on nitrogen; O_3 = ozone; POPs = moss survey on POPs.

Historically, the ICP Vegetation has focussed on the impacts of ozone pollution on vegetation. In recent years, the ICP Vegetation has reported on the widespread occurrence of ozone damage to vegetation (Hayes et al., 2007; Mills et al., 2011a), the threat of ozone pollution to food security (Mills and Harmens, 2011), impacts of ozone on the carbon sequestration in the living biomass of trees (Harmens and Mills, 2012), and impacts of ozone on ecosystem services and biodiversity (Mills et al., 2013). Furthermore, the ICP Vegetation has been instrumental in developing the methodology for establishing ozone critical levels for vegetation (Mills et al., 2011b; LRTAP Convention, 2011). The ICP Vegetation is also studying the interactive impacts of ozone and nitrogen on vegetation and the impacts of ozone on vegetation in a future climate (Vandermeiren et al., 2009). Participation from EECCA and SEE countries in the ozone work of the ICP Vegetation is very limited (Table 1.1).

Since 2000/1, the ICP Vegetation has conducted the European moss survey on heavy metals. It involves the collection of naturally-occurring mosses and determination of their heavy metal concentration at five-year intervals. European surveys have taken place every five years since 1990, and the latest survey was conducted in 2010/11. Mosses were collected at thousands of sites across Europe and their heavy metal (since 1990; Harmens et al., 2010, 2013c), nitrogen (since 2005; Harmens et al., 2011, 2013c) and POPs concentration (pilot study in 2010; see Harmens et al., 2013b) were determined. Participation from a limited number of EECCA countries mainly concerns the European moss survey on heavy metals, whereas participation in the moss survey in SEE countries is more widespread and also included nitrogen in a limited number of countries (Table 1.1).

1.3 Aim and structure of the report

In this report we review the current knowledge on the deposition of air pollutants to and their impacts on vegetation in EECCA and SEE countries (Chapter 2) and South-East Asia (Chapter 3). Knowledge is presented for the pollutants nitrogen, ozone, heavy metals, POPs and aerosols including black carbon (South-East Asia only). Conclusions and recommendations are presented in Chapter 4. Some countries have provided a country report and these are included in Annex 1; the annex also contains a country report submitted by Egypt.

2. Air pollution deposition and impacts on vegetation in EECCA and SEE

Harry Harmens, Gina Mills, Katrina Sharps, with contributions from Max Posch, Jaap Slootweg, Jean-Paul Hettelingh, Ilia Ilyin, Michael Gauss, Anna Benedictow

2.1 Introduction

Each year EMEP (European Monitoring and Evaluation Programme; <u>http://www.emep.int</u>) provides scientific information on transboundary air pollution fluxes inside the EMEP area, relying on information on emission sources and monitoring results provided by the Parties to the LRTAP Convention. EMEP consists of three main elements: (1) collection of emission data, (2) measurements of air and precipitation quality and (3) modelling of atmospheric transport and deposition of air pollutions. Officially submitted emission data by Parties to the Convention are used as input to atmospheric chemistry and transport models developed by EMEP. The Meteorological Synthesizing Centre - West (MSC-W) is responsible for the modelling assessment of sulphur, nitrogen, photo-oxidant pollutants and atmospheric particles. The modelling development for heavy metals and POPs is the responsibility of the Meteorological Synthesizing Centre - East (MSC-E). The performance of the models is evaluated through field-based measurements of air concentrations and deposition within the EMEP monitoring network.



Figure 2.1 EMEP monitoring network for acidifying, eutrophying compounds and particulate matter (PM; top left; excluding ozone only measurements sites), ozone (top right), heavy metals (bottom, left; note: Cyprus is misplaced in the map to fit inside the map) and persistent organic pollutants (POPs; bottom, right) operational in 2011. Sources: Hjellbrekke and Fjæraa (2013); Hjellbrekke et al. (2013); Aas and Breivik (2013).

Figure **2.1** and **Table 2.1** provide an overview of the EMEP measurement sites in operation in 2011. Not all sites measure all pollutants and the number of monitoring sites is limited, particularly in EECCA but also SEE countries compared to the rest of Europe. For heavy metals, 68 sites were measuring cadmium and/or lead concentrations in air, precipitation or both in 2011, of which only half were measuring one form mercury (Aas and Breivik, 2013).

Country	Station	Latitude	Longitude	A&E compounds & PM*	Ozone	Heavy metals	POPs
EECCA							
Armenia	Amberd	40 23 4 N	44 15 38 E	Х	Х		
Belarus	Vysokoe	52 20 0 N	23 26 0 E	Х			
Georgia	Abastumani	41 45 18 N	42 49 31 E	Х			
Moldova	Leova II	46 29 18 N	28 17 0 E	Х			
Russian Federation	Janiskoski	68 56 0 N	28 51 0 E	Х			
Russian Federation	Pinega	64 42 0 N	43 24 0 E	Х			
Russian Federation	Danki	54 54 0 N	37 48 0 E	Х			
Russian Federation	Lesnoy	56 31 48 N	32 56 24 E	Х			
SEE country							
Bulgaria	Rojen peak	41 41 45 N	24 44 19 E		Х		
Croatia	Puntijarka	45 54 0 N	15 58 0 E	Х			
Croatia	Zavizan	44 49 0 N	14 59 0 E	Х			
Cyprus	Ayia Marina	35 2 20 N	33 3 29 E	Х	Х	X ^{Hg}	Х
Greece	Aliartos	38 22 0 N	23 5 0 E		Х		
Greece	Finokalia	35 19 0 N	25 40 0 E		Х		
Macedonia	Lazaropole	41 19 12 N	20 25 12 E	Х	Х		
Montenegro	Zabljak	43 9 0 N	19 8 0 E	Х			
Romania	Poiana Stampei	47 19 29 N	25 8 4 E	Х	х	х	
Serbia	Kamenicki vis	43 24 0 N	21 57 0 E	Х		х	
Slovenia	lskrba	45 34 0 N	14 52 0 E	Х	х	X ^{Hg}	Х
Slovenia	Zarodnje	46 25 43 N	15 0 12 E		Х		
Slovenia	Krvavec	46 17 58 N	14 32 19 E	Х	Х		
Slovenia	Kovk	46 7 43 N	15 6 50 E		х		

 Table 2.1
 EMEP monitoring sites operational in 2011 in EECCA and SEE countries, the components measured at each site are indicated.

* Acidifying (A), eutrophying (E) compounds and particulate matter (PM); X^{Hg}: also measuring mercury (Hg).

The Working Group on Effects (http://www.unece.org/env/lrtap/workinggroups/wge/welcome.html) provides scientific support to the LRTAP Convention by developing dose-response relationships for effects of air pollutants on ecosystems, human health and materials. Based on these relationships, critical loads and levels are calculated to protect the environment and human health from adverse impacts of air pollutants (LRTAP Convention, 2011). A critical load or level is defined as a quantitative estimate of an exposure to one or more pollutants below which significant harmful effects on specified sensitive elements of the environment do not occur according to present knowledge (Nilsson and Grennfelt, 1988). The main objective of EMEP and WGE is to provide governments and subsidiary bodies under the LRTAP Convention with scientific information to support the development and further evaluation of the international protocols on emission reductions negotiated within the Convention. In the following sections we will describe the deposition of air pollutants to and impacts on terrestrial ecosystems for nitrogen (N), ozone (O₃), heavy metals and POPs.

2.2 Nitrogen

2.2.1 Background

Nearly 80% of the earth's atmosphere is made up of di-nitrogen (N_2). Whereas N_2 is unreactive and cannot be assimilated by most organisms, many reactive nitrogen (N_r) forms are essential for life, but are naturally in short supply. These include ammonia, nitrates, amino acids, proteins and many other forms (Sutton et al., 2011). At the start of the twentieth century, several industrial processes were developed to fix N_2 into N_r . Since the 1950s, the production of N_r has greatly increased to meet the demand for fertilizer to feed the world's growing population as well as industrial and other needs for N_r . Emission sources of reduced forms of N are primarily related to agricultural activities such as animal husbandry (manure) and the application and production of fertilizers. The main anthropogenic sources for oxidised forms of N are combustion processes in transport, industry and energy

production (up to 70% of oxidised N emissions). Oxidised N forms also contribute to the formation of ozone (O_3). However, N_r is a key societal threat to the environment, with adverse effects on water quality, air quality, greenhouse balance, ecosystems and biodiversity, and soil quality. Cost-benefit analysis highlights how the overall environmental costs of all N_r losses in Europe (70 – 320 billion Euros per year) outweigh the direct economic benefits of N_r in agriculture (Sutton et al., 2011).

2.2.2 Nitrogen critical load exceedances

N_r has been identified as one of the major drivers of biodiversity loss in Europe, in particular for vegetation diversity. Impacts on vegetation diversity are through direct foliar damage, eutrophication (N enrichment), acidification, and susceptibility to secondary stress (Dise et al., 2011). Evidence is strong that plant communities respond to the accumulated pool of plant-available N in the soil, therefore, the cumulative load of N to ecosystems is probably highly important. European emission controls for sulphur and N are based on the critical loads concept, an effects-based approach (Spranger et al., 2008; Bobbink and Hettelingh, 2011; LRTAP Convention, 2004; Reinds et al., 2008).



Figure 2.2 Areas in Eastern Europe where critical loads for eutrophication are exceeded by nutrient nitrogen deposition in 1990 (left) and 2010 (right). Note: The 1990 map is on the EMEP 50 km x 50 km grid whereas the 2010 map is on the 0.5 x 0.25 degree longitude – latitude grid using the latest EMEP model output; the critical loads are from the 2011/12 data base. Source: Coordination Centre for Effects, RIVM, Bilthoven, The Netherlands.

Although implementation of air pollution abatement policies developed under the LRTAP Convention have resulted in a decline of the exceedance of N critical loads across Europe since 1980, terrestrial N enrichment continues to be a serious threat to European ecosystems (Hettelingh et al., 2012; Working Group on Effects, 2013a). Areas most at risk in recent decades were those in Western and Central Europe, and those predicted to be most at risk in 2020 are parts of the Netherlands and West-France (Hettelingh et al., 2012). Figure 2.2 shows the areas at risk of eutrophication in 1990 and 2010 for EECCA and SEE countries for which data were available. The magnitude of exceedance has clearly declined in the last two decades, with areas at highest risk in 1990 being areas in the central part of the region and those nearest to Central Europe. The area on the border with Central Europe still showed the highest exceedances in 2020. The recently updated Guidance Document on Health and Environmental Improvements (Working Group on Effects, 2013b) reflects the commitments of the Revised Gothenburg Protocol to include 2005 as the base year for emission reductions reporting and 2020 emission reduction commitments. For EECCA countries, the Guidance Document only contains information for Belarus, Moldova, Russian Federation and Ukraine. The 2020 emission reduction commitments for Belarus, Moldova and Ukraine will not result in a change in the percentage of area where critical loads of N are exceededed in 2020 compared to 2005, i.e. the area of exceedance will remain 100% (Table 2.2), but the average accumulated exceedance (AAE; Posch et al., 2001), an indication of the magnitude of exceedance, will be reduced by 14 (Belarus) to 24% (Moldova).

In the Russian Federation, the percentage area of exceedance will decline from 48% in 2005 to 40% in 2020, with AAE being reduced by 33%. The predicted improvements of AAE in the EECCA

countries are relatively low compared to those in SEE countries (apart from Cyprus and Albania) and those in countries in other parts of Europe (Table 2.2; Working Group on Effects, 2013b). Although large areas in Europe are predicted to still exceed the N critical loads in 2020 under the current Gothenburg Protocol emission reduction commitments, the magnitude of exceedance as indicated by the AAE will be much lower. Hence, eutrophication will remain a threat to ecosystems in the near future, preventing recovery of ecosystems from higher N deposition in recent decades.

Table 2.2	Percentage area per country where critical loads for eutrophication of ecosystems are
	exceeded and average accumulated exceedance (AAE) in those areas for the base year 2005
	and the 2020 emission reduction commitments of the revised Gothenburg Protocol. The
	percentage improvement between 2005 and 2020 is also shown. Source: Working Group on
	Effects (2013b). Guidance document for Health and Environmental Improvements.

	20	05	20	20	% improvement 2005-2020		
	Area (%)	AAE (eq ha ⁻¹ a ⁻¹)	Area (%)	AAE (eq ha ⁻¹ a ⁻¹)	Area (%)	AAE (eq ha ⁻¹ a ⁻¹)	
EECCA							
Belarus	100	460	100	397	0	14	
Moldova	100	407	100	309	0	24	
Russian Federation	48	78	40	52	17	33	
Ukraine	100	520	100	424	0	18	
SEE							
Albania	92	289	81	218	12	25	
Bosnia & Herzegovina	72	233	67	131	7	44	
Bulgaria	77	165	38	52	51	68	
Croatia	96	502	82	262	15	48	
Cyprus	100	281	100	243	0	14	
Greece	100	377	95	219	5	42	
Macedonia	91	280	73	151	20	46	
Romania	99	493	92	269	7	45	
Slovenia	91	265	34	42	63	84	

2.2.3 Nitrogen concentrations in mosses

Several studies have shown that mosses have the potential to be indicators of atmospheric N deposition (Harmens et al., 2011, and references therein). However, sometimes the relationship between atmospheric N deposition and the N concentration in mosses is weak (e.g. Stevens et al., 2011) or shown to be species-specific (Arroniz-Crespo et al., 2008; Salemaa et al., 2008). In 2005, ectohydric moss species were sampled for the first time at the European scale to indicate spatial patterns of atmospheric N deposition across Europe (Harmens et al., 2011). Detailed statistical analysis of the European moss data (Schröder et al., 2010a) revealed that the total N concentration in mosses is significantly and best correlated with EMEP modelled air concentrations and atmospheric N deposition rates in comparison to other predictors that might contribute to the spatial variation of N concentrations in mosses. The variation in the total N concentration in mosses was best explained by the variation in ammonium (NH_4^+) concentration in air, followed by nitrogen dioxide (NO_2) concentrations in air. An apparent asymptotic relationship was found between EMEP modelled total atmospheric N deposition and the total N concentration in mosses (Harmens et al., 2011). Factors potentially affecting this relationship were discussed in more detail in the same study. Saturation appears to start at N deposition rates of ca. 15 kg ha⁻¹ y⁻¹, which might indicate the threshold of adverse impacts of N on the moss species sampled. For many habitats in Europe a N deposition of 15 kg ha⁻¹ y⁻¹ is within the range or even above the empirical critical load for N (Bobbink and Hettelingh, 2011).

So far, no EECCA countries have reported data on the N concentration in mosses and limited data is available from SEE countries. N concentrations in mosses have been determined in 2005 and 2010 in Bulgaria, Croatia, Macedonia and Slovenia (**Table 2.3**). The data for these countries are within the mid-range of those observed in European countries in general (Harmens et al., 2011; Harmens et al., 2013c). In all these countries, a decline in the N concentration in mosses has been observed between

2005 and 2010, indicating that the atmospheric deposition of N has declined since 2005 in these countries. The decline (30%) in the median concentration in mosses in Slovenia is most likely due to a change in methodology (see Annex 1), where the impact of canopy drip on N concentrations in mosses (Kluge et al., 2013) was minimised in the 2010 survey.

	N mo	ss (%)	Decline (%)
Country	2005	2010	2005-2010
Bulgaria	1.37	1.32	4
Croatia	1.60	1.49	7
Macedonia	1.21	1.06	13
Slovenia	1.84	1.29	30
Turkey	1.41	-	-

Table 2.3Median nitrogen concentration in mosses in countries in South-East Europe that reported these
concentrations for the 2005 and 2010 European moss survey.

Little is known about the relationship between N concentration in terrestrial mosses and impacts of N on terrestrial ecosystems (Harmens et al., 2012b). Empirical critical loads have been defined for various habitats (Bobbink and Hettelingh, 2011), however, the effects indicators for exceedance have not been related so far to N concentrations in mosses per se. For many terrestrial ecosystems with an empirical critical load below 15 kg ha⁻¹ y⁻¹ N effects have been reported on moss species (e.g. changes in moss species composition or abundance).

2.3 Ozone

2.3.1 Background

Ozone (O_3) is a secondary air pollutant formed, and destroyed, by a series of complex photochemical reactions involving nitrogen oxides (NO_x = NO + NO₂), methane (CH₄), carbon monoxide (CO) and non-methane volatile organic carbons (NMVOC) (Avnery et al., 2011; Royal Society, 2008). Although tropospheric (ground-level) O_3 is a natural phenomenon, since the industrial revolution O_3 concentrations in the troposphere have substantially increased from around 10-15 parts per billion (10-15 ppb), to present day values of 30-40 ppb (Simmonds et al., 2004; Sitch et al., 2007) with the steepest rise being from 1950 to 2000 (Vingarzan, 2004; Parrish et al., 2012). Future projections of O₃ concentrations are closely coupled to levels of anthropogenic precursor emissions (Dentener et al., 2006). With the global population estimated to reach 9.2 billion by 2050, associated increased demand for resources such as fossil fuels, energy production, transport and agriculture is likely to further increase precursor emissions (Oltmans et al., 2006). Tropospheric O₃ pollution is, therefore, a major concern at a local, regional and global (hemispheric) scale (Jenkin, 2008; Van Dingenen et al., 2009). Future O₃ trends will not only depend on the anthropogenic emission levels of precursors, but also on trends in temperature, humidity and solar radiation. A multi-model study of impacts of climate change alone on O_3 concentrations in Europe predicts increases in the mean O_3 concentration in the range 0.9 to 3.6 ppb for 2040-49 climates compared to 2000-09 climates (Langner et al., 2012).

As well as these steady increases in background O_3 concentrations across Europe, it is also of concern that O_3 episodes frequently occur in which the O_3 concentration exceeds 60 ppb, sometimes for several days at a time. In recent hot, dry years, O_3 episodes have been widespread across Europe. For example, in July 2006, two significant O_3 episodes occurred between 17 - 22 July and 25 - 28 July. During these episodes, O_3 concentrations in excess of 90 ppb were experienced in countries such as the UK, Belgium, Netherlands, France, Germany, Switzerland and Italy with the highest one hour value recorded being over 180 ppb in Italy (EEA, 2007). Even in a cooler, wetter year such as 2011, the EU's information threshold (one hour at 180 µg m⁻³ (or 90 ppb) was exceeded in 16 Member States whilst the alert threshold of 240 µg m⁻³ (or 120 ppb) was exceeded in Bulgaria, France, Greece, Italy, Portugal and Spain. O_3 concentrations are usually highest in rural and upland areas downwind of major conurbations, where unlike in cities, fewer other pollutants are present to react with O_3 to reduce the concentration. These rural/upland areas are where many of the ecosystems occur that provide essential services for man (agricultural production, forest production, water catchments etc.). Here, O_3 impacts on ecosystems will vary from direct toxicity and cell

damage, to indirect effects mediated by changes in individual organisms and their ecological interactions, and in the rate and nature of chemical and biological processes (Ainsworth et al., 2012; Ashmore, 2005).

Excessive uptake of O_3 by vegetation in the short-term can cause altered physiology (photosynthesis, respiration, C allocation and stomatal functioning), reduced growth (both above– and below-ground), altered phenology and increased senescence (Mills et al., 2013). In the long-term, it may lead to changes in species and genetic composition and functioning of (semi-)natural plant communities (hence, ecosystems), and to changes in water economy and C stocks. Thus, O_3 acts primarily on the processes which underlie the functioning of ecological systems whilst the benefits we derive from ecosystems are often many steps removed from these functional processes (Mills et al., 2013). Recently, the ICP Vegetation has reviewed impacts of O_3 on food security (Mills and Harmens, 2011), carbon sequestration (Harmens and Mills, 2012) and ecosystem services and biodiversity (Mills et al., 2013).

The ICP Vegetation has been instrumental in developing O_3 risk methodology for application at the European scale. Initially, O_3 exposure indices based on accumulated exposure above a threshold concentration (e.g. AOT40) were recommended for use across Europe. In the last decade, a method has been developed that takes into account the instantaneous effects of climatic factors (temperature, humidity, light), soil factors (soil moisture) and plant factors (growth stage) on the amount of O_3 that is taken up by the stomatal pores on the leaf surface (O_3 flux or phytotoxic O_3 dose over a threshold flux of Y, POD_Y). O_3 effects detected in the field are better correlated with O_3 flux than AOT40 (Hayes et al., 2007; Mills et al., 2011a), and the flux-based methodology has now been accepted by the LRTAP Convention as the preferred approach within the revised Gothenburg Protocol (Mills et al., 2011b). Critical levels of O_3 for vegetation are defined in detail in Chapter 3 of the Modelling and Mapping Manual of the LRTAP Convention (<u>http://icpvegetation.ceh.ac.uk/manuals/mapping_manual.html</u>). Data used for developing critical levels of O_3 for vegetation are primarily based on data from other parts of Europe than the EECCA and SEE region, due to a lack of data from that region.

2.3.2 Field-based evidence of ozone impacts

In 2007, the ICP Vegetation reported on field-based evidence of widespread O_3 damage to vegetation in Europe for the period 1990 to 2006 (Hayes et al., 2007). Although evidence was provided for impacts in the Eastern Mediterranean area, data was limited to impacts observed at selected sites in Greece and Slovenia, with the longest time series of data being available for Slovenia. In Slovenia (Ljubljana), the three-month AOT40 during June - August was very variable from year to year and ranged from 8.1 ppm h to 18.6 ppm h during 2003 to 2006. During April - October 2003, O_3 concentrations in Ljubljana frequently exceeded 60 ppb, with several O_3 episodes exceeding 80 ppb during June - August. In Greece, during June - August 2003 and 2004, O_3 concentrations in Kalamata and Athens reached values up to 120 and 100 ppb respectively (Hayes et al., 2007). In 1998, O_3 peaks up to 180 ppb were recorded during a four-day period in October, resulting in commercial value losses of €15,000 overnight in glass houses cultivating lettuce (**Figure 2.3**). O_3 injury has also been reported for wheat, maize, cotton, potato, water melon, musk melon, bean, onion, grapevine, courgette, chicory and fodder crops (Velissariou and Davidson, 1996; Velissariou and Skretis, 1999; Velissariou et al., 2000; Fumagali et al., 2001; see Annex 1 for further details).



Figure 2.3 Ozone-induced leaf damage in lettuce (left), cotton (middle) and water melon (right) in Greece. Source: Dimitris Velissariou.

Injury records from this region are mainly from crops; there are only a few records for trees and shrubs and no records of O₃ injury on grasses or forbs. It may well be that in this region O₃ uptake is high in crops as they are often irrigated, whereas the majority of naturally occurring vegetation experiences hot, dry conditions during the highest O_3 episodes, coinciding with low stomatal conductances and thus low stomatal O_3 fluxes. When exposing plants in biomonitoring studies to ambient air, extensive O3-induced visible leaf injury has been observed on the O3-sensitive variety of white clover (Hayes et al., 2007) and bean in Greece and Slovenia and also on Bel-W3 tobacco plants exposed at several sites across the Greater Athens Region. In addition, Velissariou et al. (1992) reported O₃-like lesions in needles of Aleppo pine within a 75 km radius from Athens. Further details on the impacts of O_3 on vegetation in Greece and Slovenia are provided in the country reports in Annex 1. Few data are available on the impacts of O₃ on vegetation in EECCA countries. In recent years, Ukraine has been participating in the O₃ biomonitoring experiments of the ICP Vegetation. In 2007, impacts of O_3 on white clover were reported for Karadag, and in 2010, impacts of O_3 on bean were reported for Kiev. In 2007, no clear impact of O₃ on the sensitive cultivar of clover was observed in Karadag. In 2010, despite O₃ injury being observed on the leaves of the sensitive variety of bean, this did not affect bean yield or weight.

2.3.3 Modelled ozone risk assessment

Due to the lack of measurements in the field on the impacts of O_3 on vegetation in EECCA and SEE countries, modelling of O_3 risk is required to assess the potential risk of O_3 on vegetation in this region. AOT40 and POD_ygen (for a generic crop and deciduous tree species) were calculated with the EMEP atmospheric transport model (Simpson et al., 2012) using the parameterisations as defined in the Modelling and Mapping Manual of the LRTAP Convention (LRTAP Convention, 2011). The data was downloaded from the EMEP web site (<u>http://webdab.emep.int/Unified_Model_Results/ydata.html</u>; downloaded 14-02-2014, model version 2013) for the extended EMEP domain.

AOT40 and POD_ygen values per 50 km x 50 km grid were averaged over five years (2007 to 2011) to smooth the annual fluctuations in these values. It should be noted that the modelled data only provide an indication of risk of O_3 impacts on crops and tree species and cannot be used to calculate exceedances of critical levels as these have not been defined so far for a generic tree species; a response function for a generic crop will be evaluated shortly.

The spatial pattern of the risk of O_3 impacts on vegetation is shown in **Figure 2.4**. As often identified before for other parts of Europe, there is a clear north south gradient for AOT40, with the risk of impact increasing from north to south. The gradient mimics the increasing level of O_3 concentration from north to south. According to the AOT40 approach, sensitive crop and forest species are most at risk from O₃ impacts in SEE countries and in the southern part of the EMEP region, in particular those areas in the east (the '-stan' countries) bordering with Central (China) and South-East Asia (Afghanistan, Pakistan, India). The AOT40-based critical level for agricultural crops and forest trees of 3 and 5 ppm h respectively is exceeded in the southern part of the EMEP region. Large parts of the Russian Federation, particularly the northern and central part, are at low risk of O₃ impacts on sensitive crop and tree species. As mentioned above, the O₃ flux-based approach is biologically more relevant and correlates better with impacts on vegetation observed in the field in other parts of Europe (Hayes et al., 2007; Mills et al., 2011a). When modelling the risk of O₃ impacts on vegetation according to the stomatal flux-based approach, determining the level of O_3 entering the leaves as a measure of phytotoxic O₃ dose (POD), then the pattern of risk is different from the concentrationbased (AOT40) approach. Both for crops and trees, the area at highest risk of O_3 impact in the EECCA countries is the region bordering with Central Europe, i.e. the western part of the Russian Federation, Belarus, Ukraine and Moldova. Many parts of SEE are also at high risk from O₃ impacts on vegetation, particularly for trees, with the risk being lowest in Turkey. It should be noted that the AOT40 and flux-based approach used here provides an estimation of the worst case for damage with adequate water supply (either rain-fed or irrigated). Reductions in ozone flux associated with dry soils such as those found in arid regions are not included in this model and thus effects may be overestimated particularly in the southern part of the region, including areas where crop irrigation is not used. More field-based evidence is required to confirm the spatial pattern of O_3 risk as indicated by the flux-based approach, as was done before for other parts of Europe (Hayes et al., 2007).





Indication of the spatial pattern of vegetation at risk of adverse impacts of ozone in EECCA and SEE countries, averaged over the period 2007-2011. Maps are shown for the concentrationbased approach (AOT40; top) and the flux-based approach (POD_Y; bottom). The flux-based risk is shown for a generic crop species (left) and a generic tree species (right) as defined for integrated assessment modelling (LRTAP Convention, 2011). Data per 50 km x 50 km grid were downloaded from the EMEP/MSC-West web site.

Note: The flux model used to generate the data in this figure provides an estimation of the worst case for damage with adequate water supply (either rain-fed or irrigated). Reductions in ozone flux associated with dry soils such as those found in arid regions are not included in this model and thus effects may be over-estimated, for example in areas where crop irrigation is not used.

2.4 Heavy metals

2.4.1 Background

In recent decades, heavy metal pollution within the EMEP region has been reduced significantly for many metals (Harmens et al., 2010, 2013c; Ilyin et al., 2013), partly due to international cooperation for pollution abatement within the LRTAP Convention. However, significant heavy metal pollution still remain in different parts of the EMEP region, particularly in south-eastern and eastern parts of Europe. Currently only two EECCA countries (Armenia and the Ukraine) have signed, and one country (Republic of Moldova) has ratified the Protocol on Heavy Metals. Little information is available on heavy metal contamination in EECCA countries and there is a need for a better coverage of heavy metal measurements, especially those for mercury in the southern and eastern parts of Europe and

Central Asia (Ilyin et al., 2013; see Figure 2.1 and Table 2.1). This might be partly achieved by involvement of national monitoring networks available in the countries. Only two countries - Republic of Moldova and Belarus officially reported their long-term trends on emissions of the three metals for 1990 – 2011 (Ilyin et al., 2013). As for other countries, their data on trends are incomplete. To fill the gaps in the emission data in these countries expert estimates are used. A peculiarity of the Caucasus countries is the fact that the contribution of non-EMEP sources to pollution is much higher than that in the western part of the EECCA region. Therefore, information about emissions in nearby non-EMEP countries presented in the EMEP domain (e.g., Iran, Syria, Iraq etc.) is needed to improve calculations of pollution levels in the EECCA countries. It should be stressed that the quality of the pollution assessment strongly depends on the availability and quality of input information, first of all, on emissions. Since monitoring information for the EECCA countries is limited, assessment of pollution levels in these countries relies entirely on modelling. Given the large gaps of knowledge on national emissions and a lack of monitoring data, assessment of pollution levels in this region is rather uncertain. Official data on anthropogenic emissions are reported only by five of the 12 EECCA countries. However, no monitoring data on heavy metal concentration in air and precipitation are reported so far. Therefore, additional efforts are needed to facilitate development of national emission inventories and monitoring networks in these countries (Ilvin et al., 2013).

2.4.2 Modelled heavy metal deposition

As for the rest of Europe, in most EECCA and SEE countries heavy metal deposition has declined between 1990 and 2011 for cadmium (Cd), lead (Pb) and mercury (Hg), with the highest reductions being computed for Pb (**Figure 2.5**). The spatial distribution of the changes in heavy metal deposition between 1990 and 2011 is shown in **Figure 2.6**. The comparable level of pollution reduction is only a characteristic of the western part of the EECCA region (Ukraine, Republic of Moldova, Belarus, Russian Federation), which is largely affected by emission changes in other European countries (Ilyin et al., 2013). Although similar reductions were computed for Pb for Kazakhstan, Uzbekistan, Turkmenistan, Kyrgyzstan and Tajikistan as for other European countries, lower reductions were computed for Cd and Hg. The lowest reductions were also reported for the eastern part of the region, i.e. Greece, Turkey and Cyprus, with increases being computed for Cd in Turkey and Cyprus (Figures 2.5 & 2.6). On average, reductions (not weighted by area) in deposition for Pb, Cd and Hg (ca. 60, 26 and 27% respectively) were similar in EECCA and SEE countries, but lower than in the rest of Europe (ca. 74, 50 and 47%).

Hg differs from other heavy metals by its long-range dispersion in the atmosphere. Therefore, Hg pollution levels in Europe are largely affected by emission sources from other regions. The major external contributor to Hg anthropogenic deposition in Europe is East Asia. Transport from this region to Europe is almost double of the reverse transport from Europe to the region. As to other regions including North and South Americas, South and Central Asia, Africa, Australia and Oceania, Europe acts as a net exporter of atmospheric mercury transporting significantly more pollution to these regions than receiving from them (Ilyin et al., 2013).





In 2011, the highest levels of metal deposition were computed in SEE countries (although lower depositions were noted for parts of Turkey and in Cyprus), the south-western part of the EECCA region (the Ukraine, Belarus, Caucasus), some eastern parts of the Russian Federation and the south-eastern (Kyrgyzstan, Tajikistan) parts of the EECCA region (Figure 2.6). Relatively high levels in these areas are caused by location of emission sources, and partly because of transboundary transport from neighbouring countries. Elevated levels of Pb and Cd deposition in 2011 are partly explained by a significant contribution of dust re-suspension. The lowest levels were found in the Arctic regions of the Russian Federation, where emissions are low. In addition, deposition of metals is low in the desert areas of Central Asia, which can be explained by low precipitation (Ilyin et al., 2013). Ilyin et al. (2013) concluded that the contributions of anthropogenic, secondary and non-EMEP sources to pollution levels differ largely among the EECCA countries. Individual peculiarities of each EECCA country should be taken into account when assessing heavy metal pollution levels.





Deposition fields of cadmium, lead and net flux of mercury in EECCA and SEE countries (top) and percentage change in deposition fields of cadmium, lead and net flux of mercury in EECCA and SEE countries between 1990 and 2011(bottom), with positive values indicating an increase and negative values a decrease. Source: EMEP/MSC-East (modified from Ilyin et al., 2013).

2.4.3 Concentrations of heavy metals in mosses

Mosses have successfully been used as biomonitors of atmospheric heavy metal deposition, in particular for Cd and Pb (Harmens et al., 2010, 2012a; Schröder et al., 2010b). The first European moss survey was conducted in 1990 and has since then be repeated at five-yearly intervals, and the latest survey was conducted in 2010 (Harmens et al., 2013c). Heavy metal concentrations in mosses provide an indication of areas at risk from high deposition of these pollutants. Temporal trends in Cd, Pb and Hg concentrations in mosses agree very well with temporal trends in their deposition modelled by EMEP/MSC-East (Harmens et al., 2010, 2012a, 2013c). One of the advantages of determining heavy metal concentrations in mosses is that it is cheaper than long-term monitoring of precipitation, so a higher sampling density can be achieved. There is a lack of EMEP monitoring stations that measure heavy metal deposition in south-eastern and eastern parts of Europe (see Figure 2.1 and Table 2.1; Ilyin et al., 2013). Some EECCA countries (or regions of EECCA countries) and several SEE countries have been participating in the European moss survey since 1990 (**Table 2.4**). **Table 2.5** provides an overview of the median concentration of heavy metals in mosses since 1995.

 Table 2.4
 Participation of EECCA and SEE countries in the European heavy metals in mosses survey since 1990. Values indicate the number of sampling sites.

EECCA	1990	1995	2000	2005	2010
Belarus		45		58	76
Russian Federation	216	621	319	220	91
Ukraine		75	115	53	17
SEE					
Albania ¹					61
Bosnia & Herzegovina			23		
Bulgaria		215	217	213	129
Croatia				94	121
Macedonia			73	72	72
Romania	56	84	214		333
Serbia			92	193	
Slovenia		29	82	57	102
Turkey				74	

¹ Data were also reported for Kosovo from 25 sites.

 Table 2.5
 Median concentration of metals in mosses sampled in EECCA and SEE countries between 1995 and 2010.

Metal		EECCA			SEE									
(mg kg ⁻¹)	Year	Belarus R	Rus. Fed.	Ukraine	Albania	Kosovo E	Bos.&Herz.	Bulgaria	Croatia	Macedonia	Romania	Serbia	Slovenia	Turkey
Arsenic	1995		0.24	0.10							0.96		0.38	
	2000		0.21	0.24			1.01	0.21		0.80	1.56	1.44	0.33	
	2005	0.15	0.23	0.22					0.37	0.68		1.41	0.43	1.71
	2010	0.12	0.15		0.24			0.63	0.28	0.69	0.68		0.26	
Cadmium	1995	0.29	0.18	0.18				0.38			0.60		0.73	
	2000		0.25	0.29				0.38		0.16	0.46		0.43	
	2005		0.24	0.32				0.31	0.28	0.29		0.26	0.33	0.30
	2010		0.07	0.36	0.11	0.13		0.21	0.38	0.22	(1.20)		0.27	
Chromium	1995	1.53	1.27	1.70				2.30			9.15		4.29	
	2000		1.43	1.50			3.45	2.41		7.46	8.46	5.07	2.59	
	2005	1.20	3.64	1.86				2.43	2.75	6.79		6.44	2.14	4.41
	2010	3.21	(9.16)	0.73	4.83	2.63		2.06	1.94	3.48	4.97		1.56	
Copper	1995	4.50	7.12	6.20				14.70			11.30		8.40	
	2000		5.84	7.31				14.51			10.82			
	2005		8.94	7.20				10.72	7.54	6.65		11.11	8.17	6.32
	2010		7.22	(21.2)	3.96	3.04		7.01	6.06	3.54	(17.79)		5.42	
Iron	1995	651	436	333				1587			1937		1007	
	2000		537	313			1350	1412		2412	2518	2365	713	
	2005	394	679	450				1399	991	2239		2267	943	1709
	2010	416	419	(1414)	1629	312		1101	789	1490	1670		548	
Lead	1995	8.16	4.45	3.40				19.0			26.5		8.55	
	2000		6.62	6.80				18.9		5.97	14.4			
	2005			7.65				14.8	2.57	7.62		16.7	10.1	5.09
	2010			7.07	2.42	7.78		8.00	3.21	4.61	(30.8)		5.01	
Mercury	1995		0.050	0.060										
-	2000		0.040	0.039						0.056				
	2005								0.064	0.068			0.095	
	2010				(0.13)	(0.033)			0.043	0.093			0.050	
Nickel	1995	1.95	4.98	2.69				3.06			2.19		2.76	
	2000		2.01	2.06			4.85	3.33		2.39	3.35	5.65		
	2005	1.25	2.74	1.70				2.99	2.68	5.82		4.43	2.75	4.04
	2010	(0.23)	2.82	(6.70)	5.81	2.00		2.61	3.16	3.45	3.60		2.12	
Vanadium	1995	3.36	3.03	1.80				4.90			6.40		4.00	
	2000		2.79	1.29			7.16	4.95		6.95	7.99	9.26		
	2005	1.33	2.27	2.13				3.88	3.10	6.38		5.76	3.38	6.28
	2010	1.19	2.45	2.63	3.52			3.07	2.55	3.49	4.89		2.30	
Zinc	1995	34.7	38.0	31.0				30.5			43.9		38.8	
	2000		35.3	29.3			23.9	32.6		39.4	(79.6)	32.6	34.5	
	2005	31.3	40.1	36.2				27.9	29.0	35.6		29.0	38.6	27.5
	2010	34.1	33.6	54.9	13.8	38.5		22.2	24.8	19.9	42.3		29.0	
Aluminium	2005	758	850	625				1495	1346	3600		3946		2260
	2010	557	922	1476	1650			1245	878	1878	3150			
Antimony	2005	0.11	0.12	0.23					0.15	0.15		0.24	0.21	0.19
,	2010	0.096	0.092								0.21		0.12	

Note: separate data were provided for Albania and Kosovo in 2010; Rus. Fed. = Russian Federation, Bosn.&Herz. = Bosnia and Herzegovina. Data in brackets indicate values not corresponding with general temporal trends (or indicate contrasting levels in neighbouring areas, e.g. for Hg in Albania and Kosovo).

Figure 2.7 shows the temporal trends in the median concentration of Cd, Pb and Hg in EECCA and SEE countries since 1995, for those countries that submitted data for at least two survey years. Limited data have been reported for Hg concentrations in mosses. Temporal trends vary per country and per heavy metal. Temporal trends can be confounded by variability in the data due to:

- The use of different moss species in different survey years;
- Variation in the sampling locations between survey years (e.g. sampling in the Ukraine was limited in the 2010 moss survey to the populated, industrialised region of Donetsk);
- Variation in the analytical methods between survey years (e.g. Macedonia used a mixture of ICP-ES and AAS in 2010 compared to INAA and AAS in 2005. For some metals the concentrations determined by INAA are higher than those determined by ICP-ES (Steinnes et al., 1997; see country report Macedonia in Annex 1);
- Sampling of mosses not affected by canopy drip from trees in forests (e.g. Slovenia in 2010 compared to previous years; canopy drip general results in higher concentrations of elements in mosses, see country report for Slovenia in Annex 1).

These confounding factors and the relatively low area of moss sampling particularly in EECCA countries should be kept in mind when comparing temporal trends (Harmens et al., 2008, 2013c).



Figure 2.7 Trends in the median concentration of cadmium, lead and mercury in EECCA and SEE countries since 1995, for those countries that submitted data in at least two survey years. Note: high values for Cd and Pb reported for Romania in 2010 are not included in the figure.

Cadmium (Cd). In earlier years the Cd concentration in mosses was higher in Bulgaria, Romania and Slovenia than in the Russian Federation or the Ukraine. However, in later years similar levels were found in the EECCA and SEE countries, although a low Cd concentration was reported for the Russian Federation in 2010. For Romania very high levels were reported for Cd in 2010, which might be related to the methodology used for Cd analysis (atomic absorption spectrometry) or sampling in very polluted areas (see Annex 1 for details). Until 2005, the average median Cd concentration in mosses in EECCA and SEE countries was higher than for the rest of Europe, with the values for 1995 and 2000 being higher in SEE than EECCA countries, but by 2010 the average median value was similar across all regions of Europe. Since 1995 the value for EECCA countries has hardly changed, whereas the values for SEE and the rest of Europe have declined considerably (ca. 63%).

Lead (Pb). In 1995, the average median Pb concentration in mosses was lower in EECCA countries than SEE countries and the rest of Europe, with the value for SEE countries being twice that of the rest of Europe. Since 1995, the value for EECCA countries has hardly changed, whereas the values for SEE and the rest of Europe have declined considerably (ca. 69%). By 2010, the value for SEE countries was still being twice that of the rest of Europe.

Mercury (Hg). Not enough data is available for Hg to make comparisons between the different regions of Europe. Whereas the concentration of mercury in moss has declined in Croatia and Slovenia between 2005 and 2010, it has increased in Macedonia since 2000.

Arsenic (As). In general, As concentrations have been lower in mosses sampled in EECCA countries than SEE countries. High As concentrations have been reported for Bulgaria (2010), Macedonia, Romania and Serbia, with lower concentrations found in Croatia and Slovenia. Since 2000, the As concentration in mosses has been similar in EECCA countries and the rest of Europe. Since 1995,

the median concentration of As has declined between 19% (EECCA) and 35% (rest of Europe) on average.

Chromium (Cr). Cr concentrations in mosses have been higher in SEE than EECCA countries due to the high levels reported for Macedonia, Romania and Serbia. On average, the median values were lower in EECCA countries than in the rest of Europe in 1995 and 2000, but the opposite was true for 2005 and 2010. Whereas the average median Cr concentration for EECCA countries has risen by 31% since 1995, it has declined by ca. 48% in SEE countries and the rest of Europe.

Copper (Cu). In 1995 and 2000, Cu concentrations in mosses were higher in SEE than EECCA countries due to high concentrations being observed in Bulgaria and Romania. However, by 2005 this difference had disappeared and since 2005 the median Cu concentration in mosses has been similar in EECCA and SEE countries and the rest of Europe. Whereas the average median Cu concentration for EECCA countries has risen by 22% since 1995, it has declined by 20 and 56% in the rest of Europe and SEE countries respectively.

Nickel (Ni). Ni concentrations in mosses were quite similar in 1995 and 2010 in EECCA and SEE countries, but were higher in SEE countries in 2000 and 2005, partly due to data being reported for Serbia where relatively high Ni concentrations were found in mosses in those years. Unusually high and low Ni concentrations were reported for Ukraine and Belarus respectively in 2010, which for Ukraine can be explained by sampling mosses in a smaller, more polluted region compared to previous years. Whereas the average median Ni concentration for EECCA and SEE countries has hardly changed since 1995, it has declined by 50% in the rest of Europe.

Vanadium (V). Relatively high V concentrations in mosses have been observed in Bulgaria, Macedonia, Romania and Serbia, in particular until 2005. Therefore, on average the median V concentration in mosses has been higher in SEE than EECCA countries, with the difference being smaller in 2010. Median values in the rest of Europe were on average similar as those in EECCA countries. Since 1995, the median V concentration has declined on average by ca. 24% in EECCA countries and the rest of Europe, and by 36% in SEE countries.

Zinc (Zn). Zn concentration in mosses vary the least between countries (Harmens et al., 2010, 2013c). Small fluctuations between years resulted in higher Zn concentrations in one of the regions in one year and in another region in another year. Between 1995 and 2010, the median Zn concentration has increased on average by 18% in EECCA countries, it has declined by 18 and 22% in the rest of Europe and SEE countries respectively.

Although we have tried to compare and generalise trends between EECCA, SEE and the rest of Europe, one should bear in mind that data availability is limited, particularly for EECCA countries. In addition, considerable variations in heavy metal concentrations in mosses and temporal trends were observed in the rest of Europe. Participation of EECCA and SEE countries in future moss surveys should be stimulated considering the fact that either relatively high levels of heavy metal pollution remain in these countries and/or the decline in pollution levels has been less than in the rest of Europe.

A review of the scientific literature showed that little is known about the relationship between heavy metal concentrations in mosses and the impacts of heavy metals on terrestrial ecosystems. Toxicity effects of heavy metals on vegetation are usually limited to areas close to pollution sources, with impacts often declining exponentially with distance from the pollution source (Harmens et al., 2012b, and reference therein). However, in the European survey, mosses are not sampled close to pollution sources and hence concentrations are often too low to be associated with an impact on terrestrial ecosystems in the sampling areas. This does not mean, however, that we should not be concerned about heavy metal deposition in remote areas as metals will accumulate in the soil and might become a problem in the future if bio-available concentrations reach critical limits.

2.4.4 Critical load exceedances for heavy metals

Although deposition of heavy metals to above-ground plant parts can lead to uptake via the leaves (Harmens et al., 2005), the risk of heavy metal toxicity to terrestrial ecosystems is often expressed as a function of the free metal ion concentration in soil solution. The LRTAP Convention has developed

the critical loads approach based on established critical limits of heavy metals in soil solution (LRTAP Convention, 2004). These critical limits are based on no-observed effect concentration (NOEC), often determined for single metals in standardised laboratory conditions for specific indicator species of toxicity. Little is known about the toxicity of metal mixtures in soil solutions and hence the NOEC for metal mixtures. Exceedance of the critical loads provide an indication of the risk of adverse impacts of heavy metals on terrestrial ecosystems. Hettelingh and Sliggers (2006) concluded that available information on the metals As, Cr, Cu, Ni, Zn and selenium (Se) suggests that none of these metals achieve high enough concentrations as a result of long-range atmospheric transport and deposition, to cause adverse effects on terrestrial ecosystems. However, although the area of exceedance of the critical loads for these heavy metals is small, even small exceedances may result in effects in the future due to the accumulative nature of heavy metals in soils. These results support the focus of the 1998 Aarhus Protocol on Heavy Metals on the metals Cd, Hg and Pb.

Table 2.6

Exceedances of critical loads for cadmium (Cd), lead (Pb) and mercury (Hg) for terrestrial ecosystem effects in 2010 in eastern and south-eastern Europe. Ex. % = percentage area exceeded; AAE = average accumulated exceedance. Source: Coordination Centre for Effects, RIVM, Bilthoven (see Slootweg et al., 2010).

	EcoArea	C	Cd	Р	b	н	g
Country	(km⁻²)	Ex. %	AAE	Ex. %	AAE	Ex. %	AAE
EECCA							
Belarus	121128	0	0.00	9	0.44	100	0.12
Moldova	2227	0	0.00	27	0.48	100	0.11
Russian Federation	1393300	1	0.01	46	4.51	100	0.16
SEE							
Albania	10082	0	0.00	15	0.79	99	0.16
Bosnia-Herzegovina	30726	0	0.00	53	3.47	100	0.22
Croatia	23666	0	0.00	19	0.77	100	0.14
Greece	30989	1	0.01	20	0.61	100	0.27
Cyprus	8148	1	0.00	33	0.89	-	-
Macedonia	12068	17	0.63	45	5.42	100	0.32
Romania	89580	0	0.00	2	0.10	100	0.15
Serbia & Montenegro	43858	1	0.01	29	1.44	100	0.19
Slovenia	13538	0	0.00	10	0.17	98	0.13



Figure 2.8 Average accumulated exceedance (AAE) of critical loads for cadmium (left), lead (middle) and mercury (right) for terrestrial ecological effects in 2010 in eastern and south-eastern Europe. Source: Coordination Centre for Effects, RIVM, Bilthoven (see Slootweg et al., 2010).

For Cd, the percentage of area with critical load exceedance for terrestrial ecotoxicological effects was generally below 1% in 2010, except for Macedonia, where the area of exceedance is 17% (**Table 2.6**). Most EMEP grids with exceedance for Cd in 2010 are in the Russian Federation (**Figure 2.8**), but the percentage of area exceeded is only 1% as the ecosystem area is large in the Russian Federation. For Pb the area and magnitude of exceedances are much higher, with exceedance occurring in all the countries included here. The area of exceedance is higher than 40% in the Russian Federation, Bosnia-Herzegovina and Macedonia and these countries also had the highest average accumulated exceedance (AAE). High exceedances were observed in a large are of the Russian Federation. The lowest percentage (2%) area of exceedance for Pb was computed for Romania. Hg has the largest exceedances, with the percentage area of exceedance being 98% or

more in all countries. The lowest exceedances for Hg were computed for northern parts of the Russian Federation, widespread exceedance was computed for all the other areas. Continued effort is required to reduce the exceedances of the critical load for Pb in the Russian Federation and continued effort is required in all EECCA and SEE countries to reduce the exceedances of the critical load for Hg. As Hg is a global pollutant, efforts are required at the hemispheric scale under the Minamata Convention to reduce Hg pollution. Hardly any field-based evidence is available to validate the critical load exceedance calculations for terrestrial ecosystems. De Zwart et al. (2010) made a first attempt to estimate the loss of species due to Cd and Pb depositions in Europe. They concluded that toxicity effects of Cd and Pb are close to zero in the vast majority of ecosystems across Europe.

2.5 Persistent organic pollutants (POPs)

2.5.1 Background

Persistent organic pollutants (POPs) are organic substances which are a concern for the environment and human health as they: possess toxic characteristics; are persistent; bioaccumulate; are prone to long-range transboundary atmospheric transport and deposition; are likely to cause significant adverse human health or environmental effects near to and distant from their source (LRTAP Convention, 1998). POPs are mainly of anthropogenic origin (Breivik et al., 2006), for example from waste incineration, industrial production and application (e.g. pesticides, flame retardants, coolant fluids) and fossil fuel burning. They show weak degradability and consequently are accumulating in the environment across the globe, including in remote areas such as the polar regions. The combination of resistance to metabolism and lipophilicity ('fat-loving') means that POPs accumulate in food chains (Jones and de Voogt, 1999). Their ecotoxicity has been highlighted in aquatic (Leipe et al., 2005) and terrestrial ecosystems (e.g. Oguntimehin et al., 2008; Smith et al., 2007).

2.5.2 Modelled deposition of POPs

Due to international cooperation and measures for pollution abatement within the LRTAP Convention, pollution from POPs has substantially decreased during the past two decades. However, levels of POP concentrations in the EMEP countries still pose risk to human health and ecosystems (Gusev et al., 2013). Currently only two of the EECCA countries (Armenia and the Ukraine) have signed, and one country (Republic of Moldova) – has ratified the Protocol on POPs. Only two countries (Slovenia and Greece) in the EECCA/SEE region monitor the deposition of POPs at EMEP monitoring sites (see Table 2.1).





Model assessment indicates reduction of POP pollution in the most of the EECCA and SEE countries from 1990 to 2011 (**Figure 2.9**). In most of the studied region, the computed deposition of HCB has dropped by more than 87% since 1990 (Gusev et al., 2013; data not shown), while for polychlorinated dibenzodioxin (PCDD) and benzo[a]pyrene (B[a]P; representing polycyclic aromatic hydrocarbons - PAHs) the decline was relatively small, especially in EECCA countries. The decrease in POP pollution particularly in EECCA countries but also SEE countries does not follow the magnitude of decline in the rest of Europe. Comparable levels of pollution reduction were seen in the western part of the

region, which is mostly affected by the emission changes in the EU countries (Gusev et al., 2013). Further improvement of the evaluation of POP pollution levels in the EECCA and SEE countries requires the refinement of information on their emissions.



Figure 2.10 Deposition fields of B[a]P, PCDD and HCB in EECCA and SEE countries in 2011 Source: EMEP/MSC-East.

Polycyclic Aromatic Hydrocarbons (PAHs). Model assessment indicates that PAH pollution represented by B[a]P decreased by 30% from 1990 to 2011 in the EMEP region (Gusev et al., 2013). The highest decline was computed for the EU countries (almost 40%). Reduction of B[a]P pollution levels was more significant during the 1990s. Pollution by PAHs varies considerably in the EECCA and SEE countries (**Figure 2.10**). In 2011, the highest levels of B[a]P deposition were computed for the western part of the region. Between 1990 and 2011, the highest reductions in B[a]P deposition were reported for the south-western part of the Russian Federation and the border with the Ukraine, Armenia, Cyprus and Albania. A slight increase in the B[a]P deposition was calculated for many eastern parts of the EECCA region (Gusev et al., 2013).

Polychlorinated dibenzo-p-dioxins and dibenzofurans (PCDD/Fs). Complex assessment of PCDD/F pollution in the EMEP countries is currently hampered by the absence of regular measurements of PCDD/Fs at the EMEP monitoring sites (Gusev et al., 2013). According to model assessments, PCDD/F pollution levels in the EMEP region decreased from 1990 to 2011 by 55%. Similar to PAHs, the highest decline was indicated for the EU countries (75%). The highest deposition of PCDD/Fs in 2011 was computed for SEE countries and the south-western part of the EECCA region. Relatively low concentrations were computed for large areas of the EECCA region (Figure 2.10). For PCDD/Fs, the decrease in deposition in the EECCA and SEE region is generally higher than for PAHs (Figure 2.9).

Hexachlorobenzene (HCB). HCB is a pollutant of global dispersion with significant potential to cycling between the environment compartments (Gusev et al., 2013). Modelling of HCB global transport and accumulation in the environment, performed on the basis of the scenario of historic and contemporary HCB emissions, indicated significant decline of HCB pollution within the EMEP region from 1990 to 2011 by almost 90%. In 2011, elevated deposition of HCB can be seen in areas with historic emissions, which led to substantial accumulation of HCB in the soil. The highest levels of HCB were generally computed for the Russian Federation, especially in the north-east, and the western part of the EECCA and SEE region (Figure 2.10).

2.5.3 POPs concentrations in mosses

Although mosses can also be used as biomonitors of atmospheric POPs pollution (Harmens et al., 2013a), the concentration of polycyclic aromatic hydrocarbons (PAHs) have only been determined at selected sampling sites in Slovenia in 2010 (Harmens et al., 2013b) and not in any other EECCA or SEE country taking part in the European moss survey. PAHs concentration in mosses in Slovenia were of a similar magnitude as those found in mosses sampled in Norway and were much lower than those reported for France, Spain (Navarra region) and Switzerland.

3. Concentrations and effects of air pollutants on vegetation in South-East Asia

Lisa Emberson, Joanne Morris, Patrick Büker, Harry Harmens, Gina Mills

3.1 Background

This review provides information describing current knowledge on concentrations, deposition and vegetation impacts of a number of air pollutants identified as particularly important across the South-East Asian region. This region encompasses the eight Malé Declaration countries: Bangladesh, Bhutan, India, Iran, Maldives, Nepal, Pakistan and Sri Lanka. The pollutants investigated are ozone (O_3) ; nitrogen species (oxidised (NO_x) and reduced (NH_y) forms that impact both as toxic gases as well as via deposition leading to soil acidification); aerosols (with a special focus on black carbon (BC)) and heavy metals (with a special focus on lead (Pb), cadmium (Cd), iron (Fe), zinc (Zn), nickel (Ni) and mercury (Hg)). This report provides new data from that presented in a previous ICP Vegetation food security report (Mills and Harmens, 2011) which had collated data up to 2010, focussing on O_3 .

Since the late 1990s, emissions of air pollutants have increased rapidly in Asia due to enhanced industrialisation, which is directly linked to continued strong economic growth of about 10% in China and India. Although legislation has attempted to curb emissions from the transport and power plant sector, emissions continue to grow, with the Asian contribution to the global emissions of SO_2 and NO_x increasing from 30 and 20% at the beginning of the 1990s to 50 and 35% in 2005, respectively (WMO/UNEP, 2011). Emissions in Asia make up a substantial fraction of the global emissions for all pollutants, with China responsible for 60 to 80% of all 'North-East Asia, South-East Asia and the Pacific' emissions. A number of reviews of emissions from the Asian region (often focussing on the main polluters, India and China) have been published recently (Cofala et al., 2007; Fang et al., 2010; Klimont et al., 2009; Wu et al., 2007) and scientists are starting to make good use of remote sensed data to verify emission inventory work and models (Klimont et al., 2009).

3.2 Emission sources

The following sectors were identified as main emitters of key air pollutants:

 NO_x . Power plants and road transport. Economic growth is accompanied by strong growth in vehicle mileage and freight transport volumes with transport demand in Asia expected to increase by a factor of 4-5. The emission control legislations present in most Asian countries will limit growth in Asian NO_x emissions by no more than 45 to 50% by 2030. In South Asia, NO_x emissions are expected to increase from 6 Tg NO_2 yr⁻¹ in 2000 to 10 Tg NO_2 yr⁻¹ in 2030 according to current legislation scenarios (Cofala et al., 2007).

Volatile Organic Compounds (VOCs, including methane - CH₄). The absence of widespread mitigation policies together with the projected economic growth indicates approximately 50% higher emissions of CH₄ for 2030 compared to 2000 (Cofala et al., 2007). In South Asia, CH₄ emissions are expected to increase from 35 Tg CH₄ yr⁻¹ in 2000 to 47 Tg CH₄ yr⁻¹ in 2030 according to current legislation. Increases are expected from all sectors except from the use of solid biomass for energy purposes.

Carbon monoxide (CO). CO is emitted whenever fossil fuels and vegetation are incompletely combusted, whether in residential stoves, industrial boilers, vehicles or through biomass burning. CO emissions have been declining globally due to a decoupling from economic growth with declining use of coal and fuel wood in domestic small stoves and the use of three-way catalysts (Cofala et al., 2007). In South Asia, CO emissions are expected to decrease from 61 Tg CO yr⁻¹ in 2000 to 52 Tg CO yr⁻¹ in 2030 according to current legislation scenarios (Cofala et al., 2007).

Ammonia (NH₃). No data could be found describing NH_3 emissions specifically for South Asia (or South Asian countries). More general studies investigating across Asia suggest that animal excreta is the main source of NH_3 emissions across the region (WMO/UNEP, 2011).

BC and organic carbon (OC). The main sector currently contributing to both BC and OC emissions is biomass burning in the residential sector (Klimont et al., 2009). BC emissions are expected to remain relatively stable in South Asia with values of 0.8 Tg C yr⁻¹ in 2000 compared with 0.7 Tg C yr⁻¹ in 2030 according to current legislation scenarios (Cofala et al., 2007). This is mainly due to the increase in road transport and likely continued reliance on traditional cooking and heating appliances and practice of crop residue burning up to 2030 in this region. OC emissions are likely to decrease rather more substantially in South Asia with values of 1.9 Tg C yr⁻¹ in 2000 compared to 1.0 Tg C yr⁻¹ in 2030 according to current legislation scenarios (Cofala et al., 2007).

Sulphur dioxide (SO₂). Emissions from power plant emissions account for over 50% of all SO₂ emissions (Klimont et al., 2009). SO₂ emissions across the whole of Asia are expected to grow by a factor of 2–3 due to the large increase in coal use for power generation without adequate pollution control. In South Asia, SO₂ emissions are expected to increase from 7 Tg SO₂ yr⁻¹ in 2000 to 22 Tg SO₂ yr⁻¹ in 2030 according to current legislation scenarios (Cofala et al., 2007).

Heavy metals. A recent review by Fang et al. (2010) provides details of atmospheric metallic element pollution in Asia over the past decade. Most studies were conducted in East Asia though the contribution sources that occur most often will be relevant for the whole of Asia. The study found that the combustion of waste and fuel generates particulate matter such as Pb, Cd, and Cr; higher concentrations of Pb were often related to high vehicle emissions close to the measurement site. Calcium (Ca), magnesium (Mg) and manganese (Mn) indicate construction materials as sources, while aluminium (Al), potassium (K), tin (Ti) and Mn indicate sources from wind-blown soils. Metallic element concentrations of Cd, Mn, Ni and Zn were significantly higher at industrial sites and are attributed to the pyrometallurgical processes (Pb and Zn smelters, non-ferrous metal industries, etc.). Analysing Hg flows in India, Burger Chakraborty et al. (2013) found coal power plants to be the predominant source of mercury emitted to the air, followed by metal production, mainly Zn. Other Hg sources are consumer products (e.g. lamps, dental amalgam) and the chlor-alkali industry.

3.3 Atmospheric concentration and deposition

This section provides some details of either monitored or modelled data of pollutant concentrations $(O_3, N \text{ deposition}, \text{ atmospheric aerosol and heavy metals})$ in South-East Asia and gives an indication of the magnitude and extent of pollutant concentrations across the region, together with the level of risk from each pollutant region-wide.

3.3.1 Ozone

Despite being a main polluter, monitored data from rural locations are relatively sparse across India and are often measured by individuals (i.e. not part of a co-ordinated monitoring campaign). A number of recent modelling studies have reviewed these *ad hoc* monitoring data in an effort to assess the accuracy of photochemical models used in risk assessment. These comparisons tend to focus on seasonal variation (Avnery et al., 2011; Van Dingenen et al., 2009) though one study also looked at diurnal variation (Engardt, 2008) in terms of capturing profiles of O_3 concentrations at locations across the region. An example of such comparisons is provided in **Figure 3.1**.

Most of these studies conclude that i) the seasonal O_3 profile is captured reasonably well; ii) O_3 concentrations tend to be overestimated, especially in Northern India. For example, Avnery et al. (2011) reported that the O_3 concentrations simulated by MOZART-2 in northern India were significantly overestimated by ca. 10-18 % throughout the year. The greater overestimates in North India may be due to the use of measured urban O_3 data for these comparisons that will tend to be lower than concentrations in rural areas. Other reasons may be due to modelled data often representing a specific height above the ground surface, e.g. Van Dingenen et al. (2009) used modelled data at a height of 30 and 10 m for the comparison, which is likely to overestimate concentrations compared to crop height. The data from Engardt (2008) shows the model tendency to

overestimate night time O_3 concentrations which may be a consequence of models struggling to represent stable conditions when O_3 would be expected to be titrated out of the atmosphere by reactions with NO_x or via deposition. Similar night time overestimations were also found in modelled vs. observed comparisons performed by Mittal et al. (2007) with the HANK model.



Figure 3.1 Comparisons of annual monthly mean surface ozone concentrations with modelled estimates based on the TM5 model for North India and South India (Reproduced from Van Dingenen et al., 2009). Dots and error bars indicate observed data, solid lines modelled data at a height of 30 m and dashed lines modelled data at a height of 10 m.



Figure 3.2 Modelled monthly mean ozone concentrations for February to May 2000 across the south Asian region using the HANK model (N.B. values given in ppb v; reproduced from Mittal et al., 2007).

In spite of these model limitations, the picture provided is of peak diurnal O_3 concentrations around midday often extending into late afternoon and during the months of October through to April. The

depression in seasonal O_3 concentrations during the monsoon period (around July to September) is also captured reasonably well in most models. Despite dissimilarities of, for example, the MATCH model (Engardt, 2008) and the HANK model (Mittal et al., 2007) regarding the absolute O_3 concentrations per region, both models show that the spatial pattern differs seasonally and that Northern India (and the corresponding South Asian region) tend to show higher O_3 concentrations than in South India (**Figure 3.2**), most likely due to a higher density of industrial and urban emissions in the north and ocean influenced atmospheric circulation patterns in the south. Hence, the models provide a useful indication of the geo-spatial extent of O_3 concentrations across the South Asian region.



Figure 3.3 Spatial distribution of (a) 90 days AOT40, (b) 75 days AOT40, (c) POD₆, and (d) POD₁₂ over the study domain for the years 2000 (left) and 2020 (right; reproduced from Tang et al., 2013).

Recently, studies have begun to focus on estimating metrics that relate O_3 concentrations to plant damage. Engardt (2008) and Mittal et al. (2007) both used the AOT40 index to characterize O_3 concentrations across the region. Both studies found AOT40 to exceed the critical level of 3 ppm h for crops set by the UNECE LRTAP Convention (LRTAP Convention, 2011) across many parts of India and particularly during the period March to May inclusive.

More recently, studies have also investigated the flux based metric (POD_Y; Mills et al., 2011b). Tang et al. (2013) simulated surface O_3 concentrations and evaluated O_3 -induced wheat production loss in China and India for the years 2000 and 2020 using dose–response functions based on AOT40 and POD_Y. Two O_3 dose metrics (90 days AOT40 and POD₆ - Pleijel et al., 2007) were derived from European experiments, and the other two (75 days AOT40 and POD₁₂ - Feng et al., 2012) were adapted from Asian studies (**Figure 3.3**). The AOT40 metrics indicated greater O_3 exposure in central-eastern China, the Pearl River Delta and Tibet for China, and around the northernmost regions, Indo-Gangetic Plain, and mega cities for India. With large future increases in O_3 concentrations, some areas in the Yangzi River Delta and the most northern part of India will experience more than 15 ppm h for 90 days AOT40 in 2020. The spatial distribution for POD₁₂ is similar to that of AOT40 as the effect of temperature on opening of the leaf pores is not considered (Tang et al., 2013). POD₆ is close to zero in the Tibetan plateau, but elevated in southern and central India due to stomatal closure in cold climates and opening in warm climates, respectively.

3.3.2 Nitrogen

Similar to the situation for O_3 , the Indian and south Asian region has not had a comprehensive systematic network to monitor deposition fluxes of N species. The 'Composition of the Atmospheric Deposition' (CAD) program was an effort to study wet and dry deposition in Asia focusing upon quality of data (<u>www.sei-international.org/rapidc/networks-cad.htm</u>). The CAD programme results highlighted the deposition of nitrate and ammonia at rural and urban sites in India (Kulshrestha et al., 2005). In India, most of the rainfall occurs during the monsoon period (June–September) and the remaining period is dominated by dry weather conditions. Even during monsoons, there are gaps of several days when it does not rain. Thus, the importance of dry deposition of atmospheric constituents in the Indian region should not be overlooked. Indeed, according to Kulshrestha et al. (2005), dry deposition of gaseous ammonia is more significant than its wet deposition in India.



Figure 3.4 Concentration (μeq l⁻¹) of ammonium (NH₄⁺; left) and nitrate (NO₃⁻; right) in rain in India. Data from measurements at rural and suburban (underlined) sites obtained with bulk (black) and wet only (red) collectors compared with the concentration field with the MATCH model for the year 2000 (reproduced from Kulshrestha et al., 2005).

Unfortunately, not many reports are available on dry deposition in India. Among the limited studies available, most of these consider dust fall as dry deposition without differentiating dry deposition of gases and particles. **Figure 3.4** shows the ammonium (NH_4^+) and nitrate (NO_3^-) concentration in rainwater in India in comparison with the concentration fields obtained with the Match model for the

year 2000. In an estimate based on EMEP dry deposition velocities, Singh et al. (2001) found that dry deposition of NH_4^+ was 9 times more significant than wet deposition at Agra. Wet deposition of NH_4^+ has been reported as 3.4 kg ha⁻¹ yr⁻¹ as compared with 39 kg ha⁻¹ yr⁻¹ of dry deposition.

3.3.3 Atmospheric aerosols

Rather than characterising concentrations for specific aerosol classes, e.g. BC, OC, sulfates etc., for which information – often only for short periods of time - is typically available for some South Asian cities only (e.g. Dutkiewicz et al., 2009; Rengarajan et al., 2007), here the focus will be on the aerosol optical depth (AOD) as a measure for assessing the total load of atmospheric aerosols of a particular area. As such AOD represents the sum of aerosols of different provenances (e.g. urban haze, smoke, (sand) dust, sea salt) distributed within a column of air ranging from the top of the atmosphere to the ground surface or vegetated canopy. The higher the AOD value, the higher the aerosol load; and the higher the aerosol load, the lower the visibility as well as the solar radiation reaching the ground surface. Remer et al. (2008) published a global study of modelled AOD based on daily MODIS (Moderate Resolution Imaging Spectroradiometer) aerosol products. South Asia exhibits especially high AOD values frequently exceeding a value of 0.4 during the northern summer, with an annual mean of approx. 0.34.

Location AOD (average for		Comment	Reference		
	indicated period)				
Trivandrum, SW India	0.37 (Dec. – May)	Average of multi-wavelength-	Krishna Moorthy et		
	0.30 (June - Aug)	radiometer (MWR) observations at	al. (2007)		
	0.32 (Oct Nov.)	four wavelengths			
Kathmandu, Nepal	0.41-0.62 (Dec. – Feb)	Ground-based measurements at	Di Girolamo (2004)		
Kanpur (IGP), India	0.44-0.72 (Dec. – Feb.)	500 nm			
Manora Peak,	0.02-0.12 (Dec. – Feb.)				
Himalaya, India					
Port Blair, Andaman	0.34 (Feb.)				
Islands, India					
Bangalore, India	0.26-0.28 (Dec. – Feb.)				
Anantapur, South	0.44-0.48 (winter)	Multi-wavelength-radiometer	Kumar et al. (2009)		
India	0.49-0.51 (summer)	(MWR) observations at ten			
	0.28-0.46 (monsoon)	wavelengths			
Hyderabad, India	0.65 (pre-monsoon)	Ground-based measurements at	Kaskaoutis et al.		
	0.46 (post-monsoon)	500 nm	(2009)		
Visakhapatnam (AP),	0.64 (May)	Ground-based measurements at	Niranjan et al.		
India	0.33 (Nov.)	500 nm	(2011)		
Kanpur (UP), India	0.57 (winter)	Ground-based measurements at	Singh (2004)		
	0.54 (pre-monsoon)	500 nm			
	0.66 (monsoon)				
	0.63 (post-monsoon)				
Dibrugarh (Assam),	0.45 (pre-monsoon)	Ground-based measurements at	Gogoi et al. (2009)		
India	0.19 (post-monsoon)	500 nm			
	0.31 (winter)				
Patiala (IGP), India	0.26 (March)	Multi-wavelength-radiometer	Singh et al. (2008)		
	0.36 (April)	(MWR) observations at 500 nm			
	0.58 (May)				

Table 2.4	Demented corrects	/ ملاحمة المحالم		a fan Cauth Aaia
Table 3.1	Reported aerosol o	plical deplin ((AOD) value	is for South Asia.

Table 3.1 summarises some observed AOD values (either using remote sensing techniques or ground-based measurements) from South Asia, averaged over different time steps varying from months to seasons. There are large variations in AOD values, mainly due to the seasonality of aerosol loading. It should be noted that in general ground stations measuring the loading and speciation of aerosol are sparse in the region (Li et al., 2007). The AOD values given in Table 3.1 are

in agreement with various other sources (e.g. Holben et al., 1998; Kaufman et al., 2002; Badarinath et al., 2007) that have confirmed that South Asia is a region of high AOD as compared to other regions, due to rapid growth leading to various anthropogenic aerosol sources and the adjacency to large arid areas (Streets et al., 2009). In particular the Indo-Gangetic Plain, South Asia's most important agricultural region, persistently has very high AOD values (Di Girolamo, 2004; Singh et al., 2008).

3.3.4 Heavy metals

Heavy metal deposition varies significantly according to location and season. Concentrations increase with proximity to emission sources, and vary with type of industry or land use as well as traffic density (Sharma et al., 2008a). Changes in climatic factors between summer, winter and monsoon seasons affect seasonal variations in heavy metal deposition (Khillare and Sarkar, 2012; Khillare et al., 2012; Shah et al., 2006), with concentrations of crustal metals such as Fe being higher in summer aided by dust storms, while industrial metals such as Cd, Ni, Zn and Cr are higher in winter due to being trapped beneath inversion layers. Rains in the monsoon season remove particles from the air, reducing atmospheric metal concentrations. Monitoring of dry deposition of heavy metals in the area surrounding Udaipur city, Rajasthan, India, Pandey & Pandey (2009a, 2009b) found steady increases in Cd, Cu, Ni and Zn from 1999 to 2005, but a significant decrease in Pb, suggested to be the result of the increasing use of unleaded fuel (**Figure 3.5**). In general, these values are deemed comparable to other regions in India, attributed to urban-industrial emissions. The higher Zn values are attributed to a nearby zinc smelter and mining.



Figure 3.5Temporal trends (1999 – 2005) in annual deposition of heavy metals in Udaipur, Rajasthan,
India (adapted from Pandey & Pandey (2009a). Note that Zn is drawn on the second axis.

Limit values and heavy metal concentrations in soil, air and plants in South Asia are shown in **Table 3.2**. For limit values, comparisons are shown with Canada, Europe and UK. The deposition of some or all of the heavy metals Zn, Cu, Cd, Pb and Fe onto fruit and vegetables across India has been found to exceed WHO and Indian national limits for safe consumption, during production (**Figure 3.6** - vegetables (Pandey and Pandey, 2009a); cereals (Pandey and Pandey, 2009b)), during transport to markets (Sharma et al., 2009), or while being sold at the roadside in urban areas: Berhampur City, Odisha (Adhikary, 2012); Varanasi, Uttar Pradesh (Sharma et al., 2008b)). This deposition comes predominantly from traffic and industrial emissions during production, traffic emissions during transport from farms to market, as well as urban traffic and industrial emissions while being sold.

Motal	Limit in agric	Min may	Moon(max)		Limit in	nnm	min may	min may littar
Weta	soil (ug g ⁻¹)	Dakistan	Varanasi IN		plants (ug	ppin min-	Dakistan	Pradech
	son (µg g)	(Ishag of al	(Sharma of al		a^{-1}	max		(P. Painai of
		(ISHAY et al.,	(3)(a)(a)(a)(a)(a)(a)(a)(a)(a)(a)(a)(a)(a)		B J	llidX (Ichog	(Hussail)	
		2015)	2009)		(Nagajyoti	(ISHAQ	2005)	al., 2010a)
					2010)	2013)	2005)	
	1				2010)	20137		
Cd	$1.4 (CA)^{-1}$	0 - 1.89	1.27 (2.3)		1.5	0-0.41		
	1.0 (UK) 2.6							
	J-U							
Ph	70 (CΔ)	4 23 - 16 24	64(103)		25	0 74-	0 02 - 0 2	05-93
1.5	250-500 (IN)	4.25 10.24	0.4 (10.5)		2.5	12 19	0.02 0.2	0.5 5.5
Fe	200 000 (111)	19 05-60 03				2 36-	18 23 -	206-1500
10		15.05 00.05				45.10	62.24	200 1000
Zn	200 (CA)	32.95 - 59	109 (133)		50	2.39-	0.97 - 2.24	7.8 - 60
	300-600 (IN)		. ,			31.25		
Ni	50 (CA)	4.21 - 18.01			1.5	1.05-	0.13 - 0.43	0.6 - 18.3
	75-150 (IN)					5.08		
Cr	64 (CA)	1.84-2.95			20	0.07-	0.09 - 0.94	0.93 - 12.1
						1.10		
	Limit in air (ng		(ng m⁻³)	mean(max)	mean ng	mean	mean ng	mean(max) ng
	m ⁻³) ³		Mashhad, Iran	ng m⁻³	m⁻³	ng m ⁻³	m ⁻³ Dhaka	m⁻³ Ahvaz, Iran
			(Pourkhabbaz	Islamabad	Tehran	Delhi	(Shah et	(Burger
			et al 2010)	(Shah et al.,	(Shah et	(Shah et	al., 2006)	Chakraborty
			ct all, 2010)	2006)	al., 2006)	al.,		et al., 2013)
						2006)		
Cd	5			2 (17)	-	6.7	2.51	
Pb	500		98.2 - 7.5	210 (4075)	1020	380	279	
Fe				930 (5979)	2230	5220	24800	
Zn			4.8 - 2.0	542 (2350)	327		801	
Ni	20		31.2 - 24.0	9 (157)	37	97		
Cr			49.6 - 9.3	42 (398)	48	104		
Hg		-						20.7 (48.6)
	India Air	min-max	Min-max: Uttar	(Pandey,				
	Quality	(peak)	Pradesh	2005)				
	Guidelines (µg	Varanasi	(Agrawal and					
	m [°]) annual	(Pandey and	Singh, 2000)					
	(24nr)	Pandey,						
Ozone	100(180)	1554)						
Ozone	100(180)	1/19)						
502	50(80)	15 - 79 (39 -	48-233	Avg: 15 - 86	1			
502	50(00)	238)	40 233	Peak: 27 -				
				191				
NO ₂	40(80)	19 - 59 (42 -						
- 2	- ()	159)						
PM10	60(100)	(TSP)			1			
		126-336						
		(215-1056)						
CA Cam	مرام الال المراثق اللا	د المثلم ما الأسم.	- 	•	•			

Table 3.2 Limit values and recorded heavy metal concentrations in soil, air and plants in South Asia. For limit values, comparisons are shown with Canada, Europe and UK.

CA - Canada; IN - India; UK - United Kingdom Canada: Canadian Environmental Quality Guidelines (<u>http://cegg-rcqe.ccme.ca/</u>)

² UK: <u>http://www.environment.agency.gov.uk</u>
 ³ Europe: <u>http://ec.europa.eu/environment/air/quality/standards.htm</u>
 ⁴ India: India Air Quality Guidelines (http://cpcb.nic.in/National_Ambient_Air_Quality_Standards.php)





3.4 Effects on vegetation

This section provides data describing site-specific pollutant effects on vegetation. Data were searched for using a variety of search terms that included the pollutant, the countries of the South Asian region and a reference to vegetation damage. These searches only found site-specific data for O_3 (**Table 3.3**) and heavy metal (**Table 3.4**) impacts. The tables provide details of the study reference, the pollutant in question, the site location, the land use type (e.g. field, forest, proximity to road traffic highways etc.), the year the study was performed, the plant species investigated, the pollutant concentration levels involved in the study, the plant response (damage parameter and magnitude) and any other information that was considered relevant.

3.4.1 Ozone impacts

Food security of many countries of South Asia is under threat due to the rapidly increasing population, industrialisation and economic growth. In Asia there are currently no air quality standards to protect agriculture from ground level O_3 . Knowledge on the impacts of O_3 on crops in South Asia was recently presented in Mills and Harmens (2011) and updated here. From studies mainly conducted in Europe and the USA, it is clear that many of the staple foods in the region are either sensitive or moderately sensitive to O₃, including maize, rice, soybean and wheat (Mills et al., 2007). The majority of studies conducted in South-East Asia report on the impacts on crop yield, although some studies also report on visible leaf injury. For example, Ahmad et al. (2013) reported 30-70% foliar injury to crop species (onion, potato, cotton) in Pakistan. Agrawal (2006) reported an increase in leaf injury from 8 to ca. 30% on tobacco along a transect away from the urban area of Varanasi, India. Evidence of the impacts of O₃ on crop yield in South-East Asia is mainly available from chemical protectant and air filtration and fumigation studies (Mills and Harmens, 2011). Data show that in India and Pakistan yield losses due to O₃ are frequently in the range of 10 to 20%, and occasionally more than 50% (Mills and Harmens, 2011). It should be noted that most of these studies were conducted at one site in India (Varanasi) and one site in Pakistan, close to Lahore. With O_3 concentrations predicted to rise in the future in South-East Asia, higher yield reductions are to be expected (see Section 3.3.1).

Since the Mills and Harmens (2011) report, most new experimental studies of O_3 impacts are from India and use filtration open top chamber (OTC) methods investigating a range of grain and vegetable crop species (rice, mustard seed, linseed, wheat, vegetables; Table 3.3). The results broadly agree with global modelling assessments of estimated O_3 damage to crops, showing reduced yield and seed quality with higher O_3 concentrations. Yield reductions of 20-35% have been recorded across species when comparing yield under 'clean' (filtered) air with ambient O_3 concentrations (Table 3.3). It was also found that increasing CO_2 concentrations mitigate the impact of O_3 (Singh et al., 2013; Mishra et al., 2013). A 40-50% reduction in oil production was recorded in linseed (Tripathi and Agrawal, 2013); in mustard seeds, Singh et al. (2013) recorded a 20% reduction in oil content and 8% reduction in protein, which was enhanced by higher CO_2 concentration.

Modelling-based studies to assess the extent and magnitude of O_3 risk to agriculture in Asia suggest that yield losses of 5–20% for important crops may be common in areas experiencing elevated O_3 concentrations (Mills and Harmens, 2011). Using a concentration-based approach, the highest relative yield losses for wheat are observed in India and China: present day losses for wheat are possibly up to 19% for China and 28% for India (Van Dingenen et al., 2009). The relative losses for rice is significantly higher in India (6-8%) than in the other regions (<5%). For soybean, high relative losses are found in China (11–21%). In terms of weight, wheat is by far the most affected crop: Van Dingenen et al. (2009) estimate a possible loss between 45 and 82 million metric tons globally, of which 30% occurs in India and 25% in China. Production losses for rice, maize and soybean are of the order 17-23 million metric tons globally. India and China account for 47% and 37% respectively of the rice production losses. The high losses obtained for India have to be considered with care, considering the large discrepancy between modelled and measured ozone concentrations in Southern India. Present day economic losses for China and India are estimated between \$3 and \$6 billion each. China and India each account for about 20% of the global economic damage. In many South-East Asian countries the relative yield losses are predicted to rise by 2030 (Van Dingenen et al., 2009).

The above-mentioned assessments have relied on European and North American dose–response relationships based on AOT40 and hence assumed an equivalent Asian crop response to O_3 for local cultivars, pollutant conditions and climate. However, comparison of the Asian data with European and North American dose-response relationships show that, almost without exception, Asian crops have a higher sensitivity to equivalent O_3 concentrations. Hence, Asian crop yield and economic loss assessments made using North American or similar European based dose-response relationships may underestimate the damage caused by ozone (Emberson et al., 2009). As such, there is an urgent need for co-ordinated experimental field campaigns to assess the effects of ozone across South-East Asia (and the rest of Asia) to allow the development of dose-response relationships for Asian cultivars and growing conditions leading to improved quantification of current and future impacts.

Tang et al. (2013) compared the flux-based and AOT40 based approach, using both European and Asian parameterisations (see Section 3.3.1 and Figure 3.3). They estimated relative yield loss (RYL) of wheat in 2000 to be 6.4-14.9% for China and 8.2-22.3% for India. POD₆ predicted greater RYL, especially for the warm regions of India, whereas the 90 days AOT40 gave the lowest estimates. For the future projection, all the O₃ dose metrics gave comparable estimates of an increase in RYL from 2000 to 2020 in the range 8.1-9.4% and 5.4-7.7% for China and India, respectively. The lower projected increase in RYL for India may be due to conservative estimation of the emission increase in 2020. Sensitivity tests of the model showed that the POD_Y-based estimates of RYL are highly sensitive to perturbations in the meteorological inputs, but that the estimated increase in RYL from 2000 to 2020 is much more robust. The projected increase in wheat production loss in China and India in the near future is substantially larger than the uncertainties in the estimation and indicates an urgent need for curbing the rapid increase in surface O₃ concentrations in this region.

3.4.2 Heavy metal impacts

While many heavy metals are essential nutrients to plant growth in small quantities, they are toxic at many of the higher concentrations reported in soils across South Asia (Table 3.4), inhibiting growth, causing chlorosis, and in some cases death (Nagajyoti et al., 2010). Laboratory-based growth experiments in India have shown:

- Reduced growth and increased free radical production at Zn concentrations higher than 5mM (Prasad et al., 1999);
- Similar effects for Cd with 40% reduction in chlorophyll at 100 μM concentration (Somashekaraiah et al., 1992);
- Increasing concentrations of Pb (10⁻⁶M 0.1M) caused significant reduction in percentage germination, shoot and root length, biomass production, and increasing inhibition of chlorophyll and protein synthesis; concentrations above 0.01M were lethal (Datta et al., 2009).
Observational studies of growth response to airborne heavy metal contamination found reduced chlorophyll a but often increased protein levels in lichen in India (Bajpai et al., 2010b; Majumder et al., 2012; Shukla and Upreti, 2007), and stunted growth in medicinal plants in Pakistan (Hussain et al., 2005). In sewage-sludge irrigated soils in India added Cd contamination above critical levels (25 mg kg⁻¹ soil) reduced yield by 20 - 54% in biomass produced for spinach, radish, coriander and fenugreek (Mani et al., 2012). Added Pb had marginal impact on yield. The added Cd also reduced sugar and vitamin C content of the vegetables significantly, and the leaves bioaccumulated up to 30% of the Cd concentration in the soil. Pb has marginal impact on vitamin C and sugar content, slightly increasing the sugar level under high Pb concentration. The accumulation of heavy metals in soils from air deposition affects plant growth, by disrupting the soil chemistry and microbial activities (Pandey and Pandey, 2009a, 2009b).

A number of studies investigated the role that different species could play as biomonitors. Species found useful for biomonitoring of heavy metal concentrations in South Asia include: lichens (Bajpai et al., 2010b; Saxena et al., 2008c); mosses (Saxena et al., 2008a,b,c; *Conocarpus erectus* (Gholami et al., 2013); pollen (Kalbande et al., 2007) or cypress, silky oak or bottlebrush (Gautam et al., 2005). Tolerant species that can be used as bioaccumulators to clean the air in urban centres have also been identified in the literature and include different species of plane trees (Pourkhabbaz et al., 2010) as well as *Morus alba, Fraxinus excesior, Cupressus sempervirens* and *Ligustrum ovalifolium* (Amini et al., 2011).

3.4.3 Nitrogen and sulphur impacts

Hardly any data describing site-specific effects of N deposition were found in the literature. The exception was a study that found a significant reduction in chlorophyll and sugar content in trees, correlated to levels of air pollution from a coal fired power plant in Uttar Pradesh. The main air pollutant at this location was SO_2 , but significant concentrations of NO_2 and particulate matter were also measured (Sharma and Tripathi, 2009). Maximum reductions of chlorophyll of 40-55% occurred in winter, when SO_2 and NO_2 concentrations were highest, and minimum reductions of 20-40% in the rainy season.

3.4.4 Aerosol impacts

Both direct and indirect effects of atmospheric aerosols on vegetation have been reported. Direct effects include a blocking of leaf pores (stomata) due to larger aerosols (Krajívcková and Mejstvrik, 1984; Kulshreshtha et al., 1994) and a change in the quantity and quality of solar radiation reaching the plant canopy (Steiner and Chameides, 2005). This second effect is of higher importance and consists of the scattering and/or absorption of solar radiation due to the aerosol loading as well as the alteration of cloud properties (Penner et al., 2001), both of which reduce the solar radiation and photosynthetically active radiation (PAR) reaching the plant surface (Moon et al., 2009; Mercado et al., 2009) while at the same time increasing its diffuse fraction (Twomey, 1977; Schwartz, 1996). The blocking of stomata can have contrasting effects on the photosynthetic activity of plants depending on the type of aerosols and plant species (Yamaguchi et al., 2014). However, the reduction in total solar radiation/PAR usually reduces photosynthetic activity (Mercado et al., 2009; Jing et al., 2010), while the increase in the diffuse fraction of solar radiation/PAR leads to an increased photosynthetic rate (Mercado et al., 2009; Zheng et al., 2010). Indirect effects are mainly associated with a change in the regional climate with knock-on effects on plant growth and yield. Rajeev and Ramanathan (2001) and Meehl et al. (2008) showed that aerosols might have an effect on the South Asian summer monsoon due to a change in the dynamics of the atmosphere. In addition, Takemura (2005) reported a negative effect of anthropogenic aerosols on the precipitation amount through a change in cloud formation. These findings are backed by an earlier study that showed how tropical aerosols can slow down the hydrological cycle (Satheesh and Ramanathan, 2000).

Ref	Pollutant	Location	Type of study	Species	Concentration	Response	Other comments
	(vear)				(averaging details)		
Ahmad et al., 2013	Ozone (2008)	Peshawar, Pakistan (8- 12km from city)	Observed injury and open top chamber (OTC)	onion (A. cepa), potato (Solanum tuberosum), cotton (Gossypium hirsutum)	36ppb (average of the overall mean summer concentrations of 2 sites)	30-70% foliar injury to all species	Spinach showed little/no effects of ozone - it is grown in winter when O ₃ conc is lower; more studies needed to confirm that damage is from O3
Kumari et al., 2013	Ozone + CO ₂ (Dec 2008 - Jan 2009)	Varanasi, India	ОТС	palak (Beta vulgaris L. var Allgreen)	Overall mean: 53 ppb (8hr daily range 36.5 - 65.8 ppb)	Yield reduction: NF:CF = 21.5% EO:NF = 25% EO:CF = 41.1%	Elevated CO ₂ mitigated impact of elevated O ₃
Tripathi and Agrawal, 2013 Note: no filtered treatment, so no measure of yield without O ₃ impact	Ozone + sUV- B (Dec 2007 - Mar 2008)	Varanasi, India	отс	linseed (<i>Linum</i> <i>usitatissimum</i> L., Linaceae) varieties Padmini and T- 397	Overall mean (NF) ppb: 50.3 EO (+10ppb) Mean monthly range: 27.7 ppb - 59.04	Yield reduction (# seeds plant ⁻¹): Padmini EO:NF = 20% T-397 EO:NF = 36% (also e.g. about 40-50% drop in oil content with high O ₃ treatment)	sUV-B had greater negative impact on yield than high O ₃ ; combined sUV-B + high O _s had least negative impact; Also give impact on seed and oil properties
Singh et al., 2013	Ozone and CO ₂ (Winter 2009/10 and 2010/11)	New Delhi, India	OTC CF: 80-85% less than ambient O ₃ NF: 5-10% less EO: NF+25- 35ppb O ₃ EO+CO ₂ : EO+500±50 ppm CO ₂ AC: chamberless air control	Indian mustard (<i>Brassica</i> <i>juncea</i> (L.) Czern.)	Average daily concentration for entire crop growth period in ambient air: 28ppb (2009/10) and 33 ppb (2010/11); max daily ambient 65 ppb in late Oct 2009	 O₃ decreased photosynthesis - by 17.6-28% in EO relative to NF: EO slows flowering (4 days longer), but speeds maturation (7-8 days earlier); and reduce yield 23- 26% relative to NF; CO₂ mitigates the negative impact of ambient O₃ on photosynthesis and yield - NF+CO₂ is increased almost to CF levels; oil content reduced by O₃: NF:CF 9% EO:NF 18-20% EO+CO₂:NF 8.5 - 10% 	Ozone reduces nutrient value of seeds

 Table 3.3
 Observed vegetation responses to surface ozone concentrations in South-East Asia, updated from the review included in Mills and Harmens (2011).

Ref	Pollutant (vear)	Location	Type of study	Species	Concentration (averaging details)	Response	Other comments
	(jear)					- protein content reduced by O_3 : EO:NF 7-8% NF:CF 13-15% inc CO_2 enhances protein loss - EO+CO ₂ :NF 12-17%	
Sarkar and Agrawal, 2012	Ozone (Jun - Oct 2007)	Varanasi, India	отс	Indian rice (<i>Oryza</i> <i>sativa L.,</i> cultivars Malviya dhan 36 and Shivani)	Mean monthly ppb: 41.3 - 59.9 (June: 42.7; July: 41.3; Aug: 44.7; Sept: 58.2; Oct: 59.9)	Yield reduction (weight of grains): Malviya dhan NF:CF = 14.3% EO++:NF = 28.4% EO++:CF = 38.7% Shivani NF:CF = 12.6% EO++:NF = 36.3% EO++:CF = 44.4%	Two levels of increased O ₃ tested (+10ppb and +20ppb). Yield reductions calculated for +20ppb
Singh et al., 2012	Ozone (Nov 2007 - Mar 2008)	Varanasi, India	OTC	mustard (<i>B.</i> <i>campestris L.</i>) varieties Vardan and Aashirwad	12 mean ppb: CF: 3.4 NF: 44.6 OP: 45 Monthly means in NF (Peaks): Nov: 41.6 (90), Dec: 27.7 (73), Jan: 46.3 (64), Feb: 48.6 (96), Mar: 59.04 (121)	Yield reduction (at RNPK): Vardan NF:CF = 7% Aashirwad NF:CF = 19.4%	No significant reduction at 1.5 RNPK (1.5 times the recommended NPK rate). Other growth parameters also measured
Tripathi and Agrawal, 2012 Note: no filtered treatment, so no measure of yield without O ₃ impact	Ozone (Nov 2010 - Mar 2011)	Varanasi, India	отс	Brassica campestris L. (cv.Sanjukta and Vardan)	Overall mean ambient (NF) ppb: 49.4 EO: ambient + 10ppb Daily mean ppb range: 26.3 (Jan) - 69.5 (Mar)	Yield reduction (# seeds plant ⁻¹): Sanjukta EO:NF = 41% Vardan EO:NF = 46%	Accumulated AOT40 was 7371 ppbh during whole growth period of plant.
Bhatia et al., 2011	Ozone (2007 and 2008)	Indian Agricultural Research Institute (IARI),	отс	Indian rice variety Pusa Sugandh-5 (PS- 5)	Mean monthly O3 (ppb, mean of two daily readings) CF: < 5ppb	2007: NF:CF = 14.4% dec yield EO:NF = 11.4% dec yield 2008:	

Ref	Pollutant	Location	Type of study	Species	Concentration	Response	Other comments
					(averaging details)		
	(year)						
		New Delhi			EO: 59.1±4.2 ppb	NF:CF = 17.9% dec yield	
					(2007); 69.7 ± 3.9 ppb	EO:NF = 12.3% dec yield	
					(2008)		
					NF: 26.2 ± 1.9 ppb		
					(2007); 37.2 ± 2.5 ppb		
					(2008)		
					AA:		
Rai et al., 2011	Ozone	Varanasi, India	OTC	Wheat	Overall daily mean for	O ₃ reduces net photosynthetic rate	
				(Triticum	study period (ppb): 45.1	and g _s during reproductive phase,	
				aestivum L.)	Overall daily mean for	impacting the assimilates required	
				cultivar	reproductive phase	for grain-filling, suggesting	
				M 533	(ppb): 50.2	significant negative repercussions	
					CF: 88% reduction cf NF	for grain yield	
Tiwari and	Ozone	Varanasi, India	OTC	Radish	Eight hourly mean	Radish:	Radish is more sensitive to
Agrawal, 2011				(Raphnus	concentration:	18.8% reduced Ps; 54.7% reduced	ozone stress than brinjal
-	(Jan-Mar			sativus L. var.	NF: 40.8 ppb	Cs; 20.8% reduced WUE	_
	2006)			Pusa Reshmi)		Brinjal:	
				and brinjal		32% reduced Ps; 26.7% reduced Cs;	
				(Solanum		17.6% reduced WUE	
				melongena L.			
				var. Pusa			
				hybrid-6)			

Ref	Pollutant	Location	Type of study (e.g. obser, exp (fum, filt etc.)	Species	Concentration (averaging details)	Response	Other comments
Rajesh Bajpai et al., 2010b	Al, Cr, Fe, Pb, Zn (2008)	Firoz Gandhi Unchahar National Thermal Power Plant Corporation (FGUNTPC), Raebareli, Uttar Pradesh	observation	Lichen (<i>P.</i> <i>cocoes</i>)	Min - max concentrations (μg g ⁻¹ dry weight, 15km - 5km from source): Fe: 206-1500 Al: 297 - 1630 Cr: 0.93 - 12.1 Cu: 0.9 - 9.9 Ni: 0.6 - 18.3 Pb: 0.5 - 9.3 Zn: 7.8 - 60	Reduced chlorophyll a concentration by up to 99%; up to 97.5% reduction in chlorophyll b; up to 80% reduction in total chlorophyll; protein and carotenoids increased	The lichen <i>P. cocoes</i> was found to be a good accumulator and therefore useful for biomonitoring
Mani et al., 2012	Cd and Pb	Allahabad, India	Field experiment	Spinach Radish Coriander Fenugreek	Concentrations added to soil: Cd: 25 & 50 mg kg ⁻¹ Pb: 250 & 500 mg kg ⁻¹ 1	1) Yield reduction (min-max across species): Cd: 23-54%, root growth inhibited Pb: marginal decrease 2) Vit C and sugar content: Cd (50mg kg ⁻¹): Up to 67% reduction Cd 50 + Pb 500 mg kg ⁻¹ : slightly less reduction, up to 60%; but mostly Pb (500 mg kg ⁻¹) addition increased sugar content	Cd above critical level (25mg kg ⁻¹ soil) impacts growth and nutritional quality significantly. Pb even at higher doses has much less impact; leafy vegetables can bioaccumulate up to 30% of added soil Cd in leaves, and at low Cd but high Pb up to 65% of added soil Cd; very little bioaccumulation of Pb
Hussain et al., 2005	Cd, Cr, Pb, Cu, Mn, Zn, Fe	Peshawar, Pakistan	observation	Milk thistle (Silybum marianum)	4 - 2 - 0km from source (mg kg ⁻¹) in leaves: Pb: 0.2; 0.02; 0.02 Cr: 0.09; 0.94; 0.79 Cd: not detected	32% reduced height, 10% reduced weight in plants at polluted site than 4km away (control);	Heavy metals accumulated in different amounts in various plant parts (roots, leaves, seeds, oil) and not always highest at pollution source

Table 3.4 Observed vegetation responses to heavy metals in South-East Asia.

Ref	Pollutant	Location	Type of study (e.g. obser, exp (fum, filt etc.)	Species	Concentration (averaging details)	Response	Other comments
					Fe: 18.23; 59.85; 62.24 Cu: 0.25; 0.49; 0.72 Mn: 1.9; 5.84; 5.68 Zn: 0.97; 2.24; 1.46 Ni: 0.16; 0.43; 0.13		
Mohsenzadeh et al., 2011 and Yousefi et al., 2009	Zn, Mn, Pb and Fe	Hamedan, Iran	observation	Reseda lutea and Chenopodiu m botrys L.	Mean soil concentration (mg kg ⁻¹): Zn: 2672 Mn: 827 Pb: 8490 Fe: 9299	 Abnormal pollen formation - thicker walls, irregular formation; new proteins found in roots 	Changes in polluted plants may be adaptations to toxic levels of metals
Prasad et al., 1999	Zn	Delhi, India	Growth experiment	Brassica juncea	Added concentration of zinc sulphate: 0.007, 0.05, 5 and 10 mM	Up to 0.05mM promoted growth; above 5mM significantly reduced growth, accelerated free radical generation	Increased enzyme activity towards detoxification stimulated by toxic Zn levels
Datta et al., 2009	Pb	India	Growth experiment	five wheat (<i>Triticum</i> <i>aestivum</i> L.) cultivars (HW 2045, HD 2733, KO307, KO 402 and HD 2954)	Added Pb concentration: 0.1M - 10 ⁻⁶ M	-significant reduction in % germination, shoot and root length, biomass production; Increasing inhibition of chlorophyll and protein synthesis	0.1M and 0.01M were lethal to germination and growth
Somashekaraiah et al., 1992	Cd	India	Growth experiment	Mung bean (Phaseolus vulgaris)	Added Cd ²⁺ concentration: 10, 50, 100μM	52% increase in lipoxygenase activity (indicates free radical production and tissue damage); 40% reduction in chlorophyll	

Ref	Pollutant	Location	Type of study	Species	Concentration	Response	Other comments
			(e.g. obser, exp (fum, filt etc.)		(averaging details)		
						production at 100 µM, 6DAG	
Majumder et al., 2012	Fe, Cr, Cu, Zn, and Pb	Kolkata, India	observation	Lichen <i>F. caperata (L.)</i> Hale	Mean site values (min-max) of 5 replicates per site: Fe: 1,970 to 3,981 mgkg ⁻¹ ;	Reduction in chlorophyll <i>a</i> with increase in Fe, Pb, Cu; MDA (membrane lipid peroxidation) increase with higher Cr, Fe, Pb; no significant difference in electrical conductivity (cell injury), but related to Fe, Cu, Pb; Zn not influencing any physiology	Lichen seen to be an efficient accumulator of Pb; Other lichen studies: (Shukla and Upreti, 2007)
Ishaq et al., 2013	Fe, Zn, Mn, Ni, Cr, Cd and Pb	Khyber Pakhtunkhwa, Pakistan	observation	Herbal drug sreen (<i>Albizia</i> <i>lebbeck</i>)	Min - max (ppm) in soil across sites (no clear gradient): Fe: 19.05-60.03 Zn: 32.95-59.00 Mn: 8.94-28.28 Ni: 4.21-18.01 Pb: 4.23-16.24 Cr: 1.84-2.95 Cd: 0-1.89 Min - max (ppm) in plant across sites (no clear gradient): Fe: 2.36-45.10 Zn: 2.39-31.25 Mn: 1.03-23.89 Ni: 1.05-5.08 Pb: 0.74-12.19 Cr: 0.07-1.10 Cd: 0-0.41		Fe and Zn may be soil contamination; Mn both soil and air contamination - also below critical limits; Ni likely soil contamination from wastewater; Pb from traffic emissions - max levels exceed WHO limit of 10 ppm; Cr from air dust - far exceed FDA limit of 0.12 ppm; Cd from soil and air - max far exceed WHO limit of 0.3 ppm

Ref	Pollutant	Location	Type of study (e.g. obser,	Species	Concentration (averaging details)	Response	Other comments
			etc.)				
Khillare et al., 2012	Cd, Cu, Cr, Ni, Zn, Fe and Mn (Jan, Jun, Aug 2009)	Delhi, India	observation	Radish (root veg), spinach (leaf veg), cowpea (legume vege), bottle gourd (fruit veg), bitter gourd (fruit veg) and ridge gourd (fruit veg)			significant seasonal variations in foliar metal concentrations; 42%, 68% and 33% of the total samples exceeded the permissible levels of Cd, Ni and Zn, but gourd vegetables
Pourkhabbaz et al., 2010	Pb, Zn, Ni, Co, Cr, Cu (May & Sept 2004 and 2005)	Mashad, Iran	observation	Plane trees (<i>Platanus</i> orientalis)	Means of 10 leaves per site (Urb - Rur) (μ g g ⁻¹): Pb: 4.5 - 1.9 Zn: 76.8-78.7 Ni: 4.2 - 4.4 Co: 0.3 - 0.4 Cr: 2.4 - 1.7 Cu: 13 - 14.5 Means of 4 air samples per site (Urb - Rur) (ng m ⁻³): Pb: 98.2 - 7.5 Zn: 4.8 - 2.0 Ni: 31.2 - 24.0 Co: 2.3 - 0.7 Cr: 49.6 - 9.3 Cu: 42 - 44.2	Leaf area, stomatal density, stomatal pore width and cuticle thickness all reduced in urban vs rural site by 16%, 28%, 31% and 32% respectively	Results show that plane trees can cope with pollution and therefore suitable to megacities
Pandey, 2005	HF, SO ₂ , TSP	Udaipur, Rajasthan,	Pot-cultured transplants to	Cassia fistula,	Annual 24hr average concentrations in	Significant stunting, reduced growth with higher	<i>P. guajava</i> most resilient, <i>C. fistula</i> most sensitive;

Ref	Pollutant	Location	Type of study (e.g. obser,	Species	Concentration (averaging details)	Response	Other comments
			exp (fum, filt etc.)				
	(2000-2001)	India	siteas around factory (pollutant source)	Psidium guajava and Carissa carandas	2000 (μg m ⁻³ min- max with distance from factory): HF: 0.2 - 3.5 SO ₂ : 15 - 86 TSD: 175 - 352	Emissions: e.g. up to 66% reduced height; foliar injury in 2 nd year only, 17-21% leaf area damaged; mean relative growth rate reduced	Impacts mainly from HF
Pandey and Pandey, 1994	SO ₂ , NO _s , O ₃ , TSP (Jan 1989 - Dec 1990)	Varanasi, India	Observation	Mangifera indica L. Psidium guajava L. Delonix regia Rafin Peltophorum pterocarpum Dalbergia sissoo Roxb Bougainville a spectabilis, Wild Carissa carandas L	Annual 24hr average concentrations in 1990 (μ g m ⁻³ min- max across pollution zones) SO ₂ : 15 - 79 NO _s : 19 - 59 O ₃ : 16 - 48 TSP: 126-336 2hr peak concentrations in 1990 (μ g m ⁻³ min- max across pollution zones) SO ₂ : 39 - 238 NO _s : 42 - 159 O ₃ : 30 - 149 TSP: 215-1056	Foliar injury (highest in winter): 6 - 19% across species in highest pollution zone vs 3 - 9% in lowest zone; Up to 31 % and 36% reduced leaf area and specific leaf weight (<i>C carandas</i>); reduction in chlorophyll, esp chlorophyll <i>a</i> : 35 - 58% dec in total chloroph.	<i>C carandas</i> is most sensitive, <i>Bougainvillea</i> is most tolerant

4. Conclusions and recommendations

Harry Harmens, Gina Mills

In this report we have provided a review on the current state of knowledge on the deposition of air pollutants to and impacts on vegetation in Eastern Europe, Caucasus and Central Asia (EECCA), South-East Europe (SEE) and South-East Asia. In these regions, there is generally a lack of a network of monitoring stations to assess the magnitude of air concentrations and depositions of pollutants. In addition, emission inventories are often incomplete or not reported at all for some pollutants, which makes it difficult to validate atmospheric transport models for these regions. Furthermore, there is a lack of coordinated monitoring networks to assess the impacts of air pollution on vegetation. Hence, the risk of adverse impacts on vegetation often has to be assessed using atmospheric transport models in conjunction with metrics developed to compute the risk of air pollution impacts on vegetation in Europe, such as critical loads and levels. Here we have focussed on the following air pollutants: nitrogen, ozone, heavy metals, POPs (EECCA/SEE countries) and aerosols, including black carbon as a component (South-East Asia).

4.1 Conclusions

4.1.1 Eastern Europe, Caucasus and Central Asia (EECCA)

Currently, for EECCA countries information on critical load exceedances for nutrient N is only available for a limited number of countries. Although the percentage area of exceedance is predicted not to change much between 2005 and 2020, the magnitude of exceedance is predicted to be reduced by 14 (Belarus) to 33% (Russian Federation) in 2020 compared to 2005, assuming full implementation of the revised Gothenburg Protocol. For modelling risks of adverse impacts of O_3 on vegetation, the concentration-based approach (AOT40) identifies the southern part of the EECCA region at highest risk, whereas the biologically more relevant flux-based approach (POD_Y) identifies the south-western part of the region bordering with Central Europe at highest risk. Both approaches identify the northern part of the region at lowest risk of adverse impacts from ozone pollution. In 2011, the highest levels of heavy metal deposition were computed for the south-western part (the Ukraine, Belarus, Caucasus), some eastern parts of the Russian Federation and the south-eastern (Kyrgyzstan, Tajikistan) parts of the EECCA region. As for the rest of Europe, in most EECCA countries heavy metal deposition has declined between 1990 and 2011 for Cd, Pb and Hg, with the highest reductions being computed for Pb. Reductions comparable to the rest of Europe were only computed for the western part of the EECCA region (Ukraine, Moldova, Belarus, Russian Federation) and for Pb in Kazakhstan, Uzbekistan, Turkmenistan, Kyrgyzstan and Tajikistan. Otherwise, lower reductions were reported for EECCA countries compared to the rest of Europe. Heavy metal concentrations in mosses have only been reported for Belarus, Russian Federation and Ukraine and critical loads exceedance data have only been reported for Belarus, Moldova and Russian Federation. Almost no critical load exceedance for terrestrial ecosystems effects has been reported for Cd, but for Pb exceedances were reported for more than 40% of the area of the Russian Federation and the area of exceedance was 100% for Hg in 2010. Model assessment indicated reduction of POP pollution in most of the EECCA countries from 1990 to 2011, although generally lower than in the rest of Europe; highest reductions were observed in the western part of the region. In 2011, the highest deposition for B[a]P and PCDD were computed in the south-western part of the region, whereas HCB levels were high for large parts of the Russian Federation.

4.1.2 South-East Europe (SEE)

Compared with Western and Central Europe, computed critical load exceedances for nutrient **N** have historically been lower in SEE and this was also the case in 2010. However, large areas are still predicted to be exceeded in 2020 with improvements since 2005 generally being lower than in Western and Central Europe. N concentrations in mosses were found to be intermediate to high in SEE compared to other European countries, indicating potentially a higher risk of N effects on ecosystems than computed by the critical loads. Field-based evidence is available for O_3 impacts particularly on crops in Greece and Slovenia, with many crops species showing visible leaf injury in

Greece. Both AOT40 and POD_Y are computed to be high in SEE, indicating that this area is at high risk of ozone damage to vegetation. Although **heavy metal** deposition has declined since 1990 in this region, the decline has generally been lower than for the rest of Europe. This might explain the relatively high concentrations of many heavy metals in mosses in countries in SEE Europe compared to the rest of Europe in recent years (Harmens et al., 2010; 2013c). Another explanation could be the amount of wind-re-suspension of historically deposited heavy metal in this more arid part of Europe, as indicated by high levels of iron and aluminium concentrations in mosses in this region. High critical load exceedances have been reported for Macedonia for Cd and Pb and for Bosnia-Herzegovina for Pb. Widespread exceedance of the critical load for Hg has been observed in this region, similar as for the rest of Europe. Model assessment indicated reduction of **POP** pollution in most of the SEE countries from 1990 to 2011, particularly for HCB, although generally lower than in the rest of Europe. Increases in B[a]P were computed for Serbia, Bulgaria and Greece.

4.1.3 South-East Asia (SEA; Malé Declaration countries)

Since the late 1990s, emissions of air pollutants have increased rapidly in Asia due to enhanced industrialisation, which is directly linked to continued strong economic growth of about 10% in China and India. Little information is available on N deposition and its impact on vegetation in SEA. According to Kulshrestha et al. (2005), dry deposition of gaseous ammonia is more important than its wet deposition in India. Staple food crops (maize, rice, soybean and wheat) are sensitive to moderately sensitive to O₃, threatening global food security. Recent flux-based risk assessment of O₃induced wheat yield loss show that the relative yield loss was 6.4-14.9% for China and 8.2-22.3% for India (Tang et al., 2013), with higher yield losses predicted for 2020, indicating the urgent need for curbing the rapid increase in surface O_3 concentrations in this region. Yield reductions of 20-35% have been recorded for various crop species when comparing yield in clear air with current ambient O₃ concentrations. South Asia is a region with high **aerosol** load compared to other regions, due its rapid growth and the arid climate. In particular the Indo-Gangetic Plain, South Asia's most important agricultural region, persistently has very high aerosol load, reducing visibility as well as solar radiation reaching the surface. Reduced photosynthesis might occur as a result of reduced solar radiation and larger aerosols blocking leaf pores, although the increase in diffuse radiation might have the opposite effect. The deposition of many heavy metals onto fruit and vegetables across India has been found to exceed WHO and Indian national limits for safe consumption.

4.2 Recommendations

This review highlights the lack of monitoring data regarding the deposition to and impacts of air pollutants on vegetation in EECCA/SEE countries and South-East Asia. It would be desirable to further enlarge coordinated networks to measure air concentrations and depositions of air pollutants, i.e. to extend the EMEP monitoring network particularly in the EECCA region and establish a similar network in South-East Asia, for example by extending the Acid Deposition Monitoring Network in East Asia (EANET) by including other regions and more pollutants. International Cooperative Programmes might consider stimulating the development of coordinated networks in these regions with the aim to establish widespread monitoring networks assessing the impacts of air pollutants on ecosystems. More measurement data are urgently needed to validate model outputs regarding the concentrations, deposition and associated risk for impacts of air pollutants on vegetation. The successful implementation of air pollution abatement policies in many other parts of Europe has highlighted the slower progress made with some of the air pollution abatement in the Eastern Europe, the Caucasus, Central and South-East Asia. Improvement of air quality in these regions will also benefit the rest of Europe due to a reduction in long-range transport of air pollutants and their precursors, particularly those of hemispheric nature such as ozone and mercury. Many air pollution issues are remaining in the studied areas that require urgent attention, especially in regions of fast economic and population growth, ensuring future sustainable development without significant impacts on the functionality of ecosystems, the services they provide and food production.

References

Aas, W., Breivik, K. (2013). Heavy metals and POP measurements, 2011. EMEP/CCC-Report 4/2013.

- Adhikary, S.P. (2012). Assessment of heavy metal toxicity on some vegetables and fruits sold in the streets of Berhampur City, Odisha, India. Pollution Research 31: 601-603.
- Agrawal, M. (2006) Crop loss assessment under ambient air pollution around Varanasi. Project report (30-137/2004/SR) submitted to University Grants Commission, New Delhi.

Agrawal, M. and Singh, J. (2000). Impact of coal power plant emission on the foliar elemental concentrations in plants in a low rainfall tropical region. Environmental Monitoring and Assessment 60: 261-282.

Ahmad, M.N., Büker, P. et al. (2013). Effects of ozone on crops in north-west Pakistan. Environmental Pollution 174: 244–249.

Ainsworth, E. A., Yendrek, C. R. et al. (2012). The effects of tropospheric ozone on net primary productivity and implications for climate change. Annual Review of Plant Biology, 63: 637-661.

Amini, H., Hoodaji, M. et al. (2011). Evaluation of some tree species for heavy metal biomonitoring and pollution tolerance index in Isfahan urban zone. African Journal of Biotechnology 10.

Arroniz-Crespo, M., Leake, J.R. et al. (2008). Bryophyte physiological responses to, and recovery from, long-term nitrogen deposition and phosphorus fertilisation in acidic grassland. New Phytologist 180: 864-874.

Ashmore, M. R. (2005). Assessing the future global impacts of ozone on vegetation. Plant Cell and Environment 28(8): 949-964.

Avnery, S., Mauzerall, D.L. et al. (2011). Global crop yield reductions due to surface ozone exposure: 1. Year 2000 crop production losses and economic damage. Atmospheric. Environment. 45: 2284-2296.

Badarinath, K.V.S., Kharol, S.K. (2007). Case study of a dust storm over Hyderabad area, India: Its impact on solar radiation using satellite data and ground measurements. Sci. Total Environ. 384: 316-332.

- Bajpai, R., Upreti, D.K. et al (2010a). Passive monitoring of atmospheric heavy metals in a historical city of central India by Lepraria lobificans Nyl. Environ. Monit. Assess. 166: 477-484.
- Bajpai, R., Upreti, D.K., et al. (2010b). Biodiversity, bioaccumulation and physiological changes in lichens growing in the vicinity of coal-based thermal power plant of Raebareli district, north India. J. Hazard. Mater. 174: 429–436.
- Bhatia, A., Ghosh, A. et al. (2011). Effect of elevated tropospheric ozone on methane and nitrous oxide emission from rice soil in north India. Agric. Ecosyst. Environ. 144: 21-28.
- Bobbink, R., Hettelingh, J-P. (Eds.) (2011). Review and revision of empirical critical loads and dose-response relationships. Proceedings of an international expert workshop, Noordwijkerhout, 23-25 June 2010. RIVM report 680359002, Bilthoven, The Netherlands.
- Breivik, K., Vestreng, V. et al. (2006). Atmospheric emissions of some POPs in Europe: a discussion of existing inventories and data needs. Environmental Science and Policy 9: 663 - 674.

Burger Chakraborty, L., Qureshi, A. et al. (2013). Anthropogenic Mercury Flows in India and Impacts of Emission Controls. Environ. Sci. Technol. 47: 8105-8113.

Cofala, J., Amann, M.et al. (2007). Scenarios of global anthropogenic emissions of air pollutants and methane until 2030. Atmospheric Environment 41: 8486-8499.

Datta, J.K., Ghanty, S.et al. (2009). Impact of lead on germination physiology of certain wheat cultivars (Triticum aestivum L.). J. Ecophysiol. Occup. Health 9: 145–151.

- De Zwart, D., Slootweg, J. et al. (2010). Loss of species due to cadmium and lead depositions in Europe. In: Slootweg, J., Posch, M. et al. (Eds). Progress in the modelling of critical thresholds and dynamic modelling, including impacts on vegetation in Europe. CCE Status Report 2010, Coordination Centre for Effects, RIVM, Bilthoven, The Netherlands.
- Dentener, F., Stevenson, D. et al. (2006). The global atmospheric environment for the next generation. Environmental Science & Technology 40(11): 3586-3594.

Di Girolamo, L. (2004). Analysis of Multi-angle Imaging SpectroRadiometer (MISR) aerosol optical depths over greater India during winter 2001-2004. Geophys. Res. Lett. 31.

Dise, N. B., Ashmore, M. et al. (2011) Nitrogen as a threat to European Terrestical Biodiversity, page 463-494. In: Sutton, M. A., Howard, C. M. et al. (Eds.) The European Nitrogen Assessment. Cambridge University Press, Cambridge.

Dutkiewicz, V.A., Alvi, S.et al. (2009). Black carbon aerosols in urban air in South Asia. Atmos. Environ. 43: 1737–1744.

EEA (2007). Air pollution by ozone in Europe in summer 2006. European Environment Agency Technical Report No 5/2007. Copenhagen, Denmark.

Emberson, L.D., Büker, P., et al (2009). Dose-response relationships derived in North America underestimate the effects of ozone (O₃) on crop yields in Asia. Atmospheric Environment 43: 1945-1953

Engardt, M. (2008). Modelling of near-surface ozone over South Asia. J. Atmospheric Chem. 59, 61-80.

Fang, G.-C., Huang, Y.-L. et al. (2010). Study of atmospheric metallic elements pollution in Asia during 2000–2007. J. Hazard. Mater. 180: 115 – 121.

Feng, Z., Tang, H.et al. (2012). A stomatal ozone flux-response relationship to assess ozone-induced yield loss of winter wheat in subtropical China. Environ. Pollut. 164: 16-23.

Fumagalli, I., Gimeno, B. S. et al. (2001). Evidence of ozone-induced adverse effects on crops in the Meditteranean region. Atmospheric Environment 35: 2583-2587.

Gautam, P., Blaha, U. et al. (2005). Magnetic susceptibility of dust-loaded leaves as a proxy of traffic-related heavy metal pollution in Kathmandu city, Nepal. Atmos. Environ. 39: 2201-2211.

Gholami, A., Davami, A.H.et al. (2013). Evaluation of "Conocarpus erectus" Plant as Biomonitoring of Soil and Air Pollution in Ahwaz Region. Middle-East J. Sci. Res. 13: 1319-1324.

Gogoi, M.M., Krishna Moorthy, K.et al. (2009). Climatology of columnar aerosol properties and the influence of synoptic conditions: First-time results from the northeastern region of India. J. Geophys. Res. 114.

Gusev, A., Rozovskaya, O. et al. (2013). Persistent organic pollutants in the environment. EMEP Status Report 3/2013.

Harmens, H., Foan, L. et al. (2013a). Terrestrial mosses as biomonitors of atmospheric POPs pollution: A review. Environmental Pollution 173: 245-254.

Harmens, H., Ilyin, I. et al. (2012a). Country-specific correlations across Europe between modelled atmospheric cadmium and lead deposition and concentrations in mosses. Environmental Pollution 166: 1-9.

Harmens, H., Mills, G. (2012). Ozone pollution: Impacts on carbon sequestration in Europe. Programme Coordination Centre for the ICP Vegetation, Centre for Ecology and Hydrology, Bangor, UK. ISBN 978-1-906698-31-7.

Harmens, H., Mills, G. et al. (2005). Air pollution and vegetation. ICP Vegetation annual report 2004/2005. Programme Coordination Centre for the ICP Vegetation, Centre for Ecology and Hydrology, Bangor, UK. ISBN: 1-870393-80-5.

Harmens, H., Mills, G. et al. (2012b). Air pollution and vegetation. ICP Vegetation annual report 2011/2012. Programme Coordination Centre for the ICP Vegetation, Centre for Ecology and Hydrology, Bangor, UK. ISBN: 978-1-906698-35-5.

- Harmens, H., Mills, G. et al. (2013b). Air pollution and vegetation. ICP Vegetation annual report 2012/2013. Programme Coordination Centre for the ICP Vegetation, Centre for Ecology and Hydrology, Bangor, UK. ISBN: 978-1-906698-43-0.
- Harmens, H., Norris, D.A. et al. (2010). Mosses as biomonitors of atmospheric heavy metal deposition: spatial and temporal trends in Europe. Environmental Pollution 158: 3144-3156.

Harmens, H., Norris, D. A. et al. (2011). Nitrogen concentrations in mosses indicate the spatial distribution of atmospheric nitrogen deposition in Europe. Environmental Pollution 159: 2852-2860.

Harmens, H., Norris, D. et al. (2013c). Heavy metals and nitrogen in mosses: spatial patterns in 2010/2011 and long-term temporal trends in Europe. ICP Vegetation Programme Coordination Centre, CEH Bangor, UK. ISBN: 978-1-906698-38-6.

Hayes, F., Mills, G. et al. (2007). Evidence of widespread ozone damage to vegetation in Europe (1990 - 2006). Programme

Hays, F., Mins, G. et al. (2007). Evidence of widespread ozone double dou

Hettelingh, JP, Sliggers, J (Eds.) (2006). Heavy Metal Emissions, Depositions, Critical Loads and Exceedances in Europe. Ministry of the Environment VROM DGM, Netherlands.

Hjellbrekke, A.-G., Fjæraa, A.M. (2013). Data report 2011: Acidifying and eutrophying compounds and particulate matter. EMEP/CCC-Report 2/2013.

Hjellbrekke, A.-G., Solberg, S., Fjæraa, A.M. (2013). Ozone measurements 2011. EMEP/CCC-Report 3/2013.

Holben, B.N., Eck, T.F. et al. (1998). AERONET-A federated instrument network and data archive for aerosol characterization. Remote Sens. Environ. 66: 1-16.

Hussain, I., Khan, L.et al. (2005). Effect of heavy metals on the growth and development of Silybum marianum, in various polluted areas of Peshawar, Pakistan. J. Chem. Soc. Pak. 27: 367-373.

Ilyin, I., Rozovskaya, O. et al. (2013). Heavy metal transboundary pollution of the environment. EMEP Status Report 2/2013.

Ishaq, M., Rehman, A. et al. (2013). Comparative study of heavy metals in Albizia lebbeck, collected from different environmental sites. Int. J. Pharm. Sci. Rev. Res. 20: 5-9.

Jenkin, M. E. (2008). Trends in ozone concentration distributions in the UK since 1990: Local, regional and global influences. Atmospheric Environment 42(21): 5434-5445.

Jing, X., Huang, J. et al. (2010). The effects of clouds and aerosols on net ecosystem CO<sub>2</sub> exchange over semiarid Loess Plateau of Northwest China. Atmospheric Chem. Phys. 10: 8205-8218.

Jones K.C. and De Voogt, P. (1999). Persistent organic pollutants (POPs): State of the science. Environmental Pollution 100: 209-221.

Kalbande, D.M., Dhadse, S.N. et al. (2007). Biomonitoring of heavy metals by pollen in urban environment. Environ. Monit. Assess. 138: 233-238.

Kaskaoutis, D.G., Badarinath, K.V.S., et al. (2009). Variations in the aerosol optical properties and types over the tropical urban site of Hyderabad, India. J. Geophys. Res. 114.

Kaufman, Y.J., Tanré, D. et al. (2002). A satellite view of aerosols in the climate system. Nature 419: 215-223.

Khillare, P.S., Jyethi, D.S. et al. (2012). Health risk assessment of polycyclic aromatic hydrocarbons and heavy metals via dietary intake of vegetables grown in the vicinity of thermal power plants. Food Chem. Toxicol. 50: 1642-1652.

Khillare, P.S. and Sarkar, S. (2012). Airborne inhalable metals in residential areas of Delhi, India: distribution, source apportionment and health risks. Atmospheric Pollut. Res. 3: 46-54. Klimont, Z., Cofala, J. (2009). Projections of SO2, NOx and carbonaceous aerosols emissions in Asia. Tellus Series B-Chem. Phys.

Meteorol. 61: 602-617.

Kluge, M., Pesch, R. et al. (2013). Accounting for canopy drip effects of spatiotemporal trends of concentrations of N in mosses, atmospheric N depositions and critical load exceedances: a case study from North-Western Germany. Environmental Sciences Europe 25: 26.

Krajívcková, A. and Mejstvrik, V. (1984). The effect of fly ash particles on the plugging of stomata. Environ. Pollut. Ser. Ecol. Biol. 36: 83-93.

Krishna Moorthy, K., Babu, S.S. et al. (2007). Temporal heterogeneity in aerosol characteristics and the resulting radiative impact at a tropical coastal station-Part 1: Microphysical and optical properties, in: Annales Geophysicae. European Geosciences Union, pp. 2293-2308.

Kulshreshtha, K., Farooqui, A. et al. (1994). Effect of diesel exhaust pollution on cuticular and epidermal features of Lantana camara L. and Syzygium cuminii L. (Skeels.). J Env Sci Health Part Env Sci Eng. Toxicol. 29: 301-308.

Kulshrestha, U., Granat, L. et al. (2005). Review of precipitation monitoring studies in India-a search for regional patterns. Atmos. Environ, 39: 7403–7419.

Kumar, K.R., Narasimhulu, K. et al. (2009). Temporal and spectral characteristics of aerosol optical depths in a semi-arid region of southern India. Sci. Total Environ. 407: 2673-2688.

Kumari, S., Agrawal, M. et al. (2013). Impact of elevated CO₂ and elevated O₃ on Beta vulgaris L.: Pigments, metabolites, antioxidants, growth and yield. Environ. Pollut. 174: 279-288.

Langner, J., Engardt, M. et al. (2012). A multi-model study of impacts of climate change on surface ozone in Europe. Atmospheric Chemistry and Physics 12: 10423-10440.

Leipe T., Kersten M. et al. (2005). Ecotoxicity assessment of natural attenuation effects at a historical dumping site in the western Baltic Sea. Marine Pollution Bulletin 50: 446-459.

Li, Z., Chen, H. et al. (2007). Preface to special section on East Asian Studies of Tropospheric Aerosols: An International Regional Experiment (EAST-AIRE). J. Geophys. Res. 112.

LRTAP Convention (2004). Manual on methodologies and criteria for modelling and mapping critical loads and levels and air pollution UNECE Convention on Long-range risks effects. and trends. Transboundary Air Pollution. http://wgecce.org/Publications/Mapping_Manual (18-02-2014).

LRTAP Convention (1998). Protocol to the 1979 Convention on Long-range Transboundary Air Pollution on Persistent Organic Pollutants. http://www.unece.org/env/Irtap/

LRTAP Convention (2011). Manual on methodologies and criteria for modelling and mapping critical loads and levels and air pollution risks Chapter 3: Mapping critical effects. and trends. levels for vegetation. http://icpvegetation.ceh.ac.uk/manuals/documents/Ch%203%20revised%20summer%202010.%20updated%20June%20201 1.pdf (12-03-2014).

Majumder, S., Mishra, D. et al. (2012). Physiological and chemical response of the lichen, Flavoparmelia caperata (L.) Hale, to the urban environment of Kolkata, India. Environ. Sci. Pollut. Res. 20: 3077-3085.

Sharma, B. et al. (2012). Cadmium and Lead Bioaccumulation During Growth Stages Alters Sugar and Vitamin C Content in Mani D Dietary Vegetables. Proc. Natl. Acad. Sci. India Sect. B Biol. Sci. 82: 477-488.

Meehl, G.A., Arblaster, J.M. et al. (2008). Effects of Black Carbon Aerosols on the Indian Monsoon. J. Clim. 21: 2869–2882.

Mercado, L.M., Bellouin, N. et al. (2009). Impact of changes in diffuse radiation on the global land carbon sink. Nature 458: 1014–1017. Mills G, Buse A, et al (2007). A synthesis of AOT40-based response functions and critical levels of ozone for agricultural and

horticultural crops. Atmospheric Environment 41: 2630-2643.

Mills, G., Harmens, H. (Eds.) (2011). Ozone pollution: A hidden threat to food security. ICP Vegetation Programme Coordination Centre, Centre for Ecology and Hydrology, UK. ISBN: 978-1-906698-27-0.

Mills, G., Hayes, F. et al. (2011a). Evidence of widespread effects of ozone on crops and (semi-)natural vegetation in Europe (1990 -2006) in relation to AOT40 - and flux-based risk maps. Global Change Biology 17: 592-613.

Mills, G., Pleijel, H. et al. (2011b). New stomatal flux-based critical levels for ozone effects on vegetation. Atmospheric Environment 45: 5064-5068.

Mills, G., Wagg, S. et al (2013). Ozone Pollution: Impacts on ecosystem services and biodiversity. Programme Coordination Centre for the ICP Vegetation, Centre for Ecology and Hydrology, Bangor, UK. ISBN: 978-1-906698-39-3.

Mishra, A.K., Rai, R., Agrawal, S.B. (2013). Differential response of dwarf and tall tropical wheat cultivars to elevated ozone with and without carbon dioxide enrichment: Growth, yield and grain quality. Field Crops Research 145: 21-32.

Mittal, M.L., Hess, P.G. et al. (2007). Surface ozone in the Indian region. Atmos. Environ. 41: 6572–6584.

Mohsenzadeh, F., Chehregani, A. et al. (2011). Effect of the Heavy Metals on Developmental Stages of Ovule, Pollen, and Root Proteins in Reseda lutea L. (Resedaceae). Biol. Trace Elem. Res. 143: 1777–1788.

Moon, K., Kim, J. et al. (2009). Effect of aerosol optical depth on diffuse photosynthetically active solar irradiance. Asia-Pac. J. Atmospheric Sci. 45: 307–317.

Nagajyoti, P.C., Lee, K.D. et al. (2010). Heavy metals, occurrence and toxicity for plants: a review. Environ. Chem. Lett. 8: 199–216. Nilsson, J., Grennfelt, P. (Eds.) (1988). Critical loads for sulphur and nitrogen. UNECE/Nordic Council report, Skokloster, Sweden.

Nisson, J., Grennier, P. (Eds.) (1988). Critical loads for suppur and hitrogen. U Nordic Council of Ministers, Copenhagen.

Niranjan, K., Spandana, B. et al., (2011). Measurements of aerosol intensive properties over Visakhapatnam, India for 2007. Ann. Geophys. 29: 973–985.

Oguntimehin, İ., Nakatani, N. et al. (2008). Phytotoxicities of fluoranthene and phenanthrene deposited on needle surfaces of the evergreen conifer, Japanese red pine (*Pinus densiflora* Sieb. et Zucc.). Environmental Pollution 154: 264-271.

Oltmans, S. J., Lefohn, A. S. et al. (2006). Long-term changes in tropospheric ozone. Atmospheric Environment 40(17): 3156-3173.

Pandey, J. (2005). Evaluation of air pollution phytotoxicity downwind of a phosphate fertilizer factory in India. Environ. Monit. Assess. 100: 249–266.

Pandey, J. and Pandey, U. (1994). Evaluation of air pollution phytotoxicity in a seasonally dry tropical urban environment. Environ. Monit. Assess. 33: 195–213.

Pandey, J. and Pandey, U. (2009a). Accumulation of heavy metals in dietary vegetables and cultivated soil horizon in organic farming system in relation to atmospheric deposition in a seasonally dry tropical region of India. Environ. Monit. Assess. 148: 61–74.
 Pandey, J. and Pandey, U. (2009b). Atmospheric deposition and heavy metal contamination in an organic farming system in a

seasonally dry tropical region of India. J. Sustain. Agric. 33: 361–378. Parrish, D. D., Law, K.S. et al. (2012) Long-term changes in lower tropospheric baseline ozone concentrations at northern mid-latitudes.

Atmospheric Chemistry and Physics Discussions 12: 11485-11504.

Penner, J.E. et al. (2001). Aerosols, their Direct and Indirect Effects, page 289-348. In: Houghton, J.T., et al. (Eds.), Climate Change 2001: The Scientific Basis. Cambridge University Press, Cambridge, UK.

Pleijel, H., Danielsson, H. et al. (2007). Ozone risk assessment for agricultural crops in Europe: Further development of stomatal flux and flux–response relationships for European wheat and potato. Atmos. Environ. 41: 3022–3040.

Posch, M., Hettelingh, J.-P. et al. (2001). Characterization of critical load exceedances in Europe. Water, Air and Soil Pollution 130: 1139–1144.

Pourkhabbaz, A., Rastin, N. et al. (2010). Influence of Environmental Pollution on Leaf Properties of Urban Plane Trees, Platanus orientalis L. Bull. Environ. Contam. Toxicol. 85: 251–255.

Prasad, K.V.S., Paradha Saradhi, P. et al. (1999). Concerted action of antioxidant enzymes and curtailed growth under zinc toxicity in Brassica juncea. Environ. Exp. Bot. 42: 1–10.

Rai, R., Agrawal, M. et al. (2011). Effects of ambient O3 on wheat during reproductive development: Gas exchange, photosynthetic pigments, chlorophyll fluorescence, and carbohydrates. Photosynthetica 49: 285–294.

Rajeev, K. and Ramanathan, V. (2001). Direct observations of clear-sky aerosol radiative forcing from space during the Indian Ocean Experiment. J. Geophys. Res. Atmospheres 1984–2012. 106: 17221–17235.

Reinds, G.J., Posch, M., et al. (2008). Critical loads of sulphur and nitrogen for terrestrial ecosystems in European and Northern Asia using different soil chemical criteria. Water, Air and Soil Pollution 193: 269-287.

Remer, L.A., Kleidman, R.G.et al. (2008). Global aerosol climatology from the MODIS satellite sensors. J. Geophys. Res. 113.

Rengarajan, R., Sarin, M.M. et al. (2007). Carbonaceous and inorganic species in atmospheric aerosols during wintertime over urban and high-altitude sites in North India. J. Geophys. Res. 112.

Royal-Society (2008). Ground-level ozone in the 21st century: future trends, impacts and policy implications. Science Policy Report 15/08.

Salemaa, M., Mäkipää, R. et al. (2008). Differences in the growth response of three bryophyte species to nitrogen. Environmental Pollution 152: 82-91.

Sarkar, A. and Agrawal, S.B. (2012). Evaluating the response of two high yielding Indian rice cultivars against ambient and elevated levels of ozone by using open top chambers. J. Environ. Manage. 95: S19–S24.

Satheesh, S.K. and Ramanathan, V. (2000). Large differences in tropical aerosol forcing at the top of the atmosphere and Earth's surface. Nature 405: 60-63.

Saxena, D.K., Singh, S. et al. (2008a). Atmospheric heavy metal deposition in Garhwal Hill Area (India): Estimation based on native moss analysis. Aerosol and Air Quality Research 8: 94-111.

Saxena, D.K., Singh, S. et al. (2008b). Retrospective metal data of the last 100 years deduced by moss, *Barbula sp.* from Mussoorie city, Garhwal Hills, India. Current Science 94:901-904.

Saxena, D.K., Srivastava, K.et al. (2008c). Biomonitoring of metal deposition by using moss transplant method through *Hypnum cupressiforme* (Hedw.) in Mussoorie. Journal of Environmental Biology 29: 683-688.

Schröder, W., Holy, M. et al. (2010a). First Europe-wide correlation analysis identifying factors best explaining the total nitrogen concentration in mosses. Atmospheric Environment 44: 3485-3491.

Schröder, W., Holy, M. et al. (2010b). Are cadmium, lead and mercury concentrations in mosses across Europe primarily determined by atmospheric deposition of these metals? Journal of Soils and Sediments 10: 1572-1584.

Schwartz, S.E. (1996). The Whitehouse Effect - Shortwave radiative forcing of climate by anthropogenic aerosols: An overview. J. Atmospheric Sci. 27: 359–382.

Shah, M.H., Shaheen, N.et al. (2006). Characterization, Source Identification and Apportionment of Selected Metals in TSP in an Urban Atmosphere. Environ. Monit. Assess. 114: 573–587.

Sharma, A.P. and Tripathi, B.D. (2009). Biochemical responses in tree foliage exposed to coal-fired power plant emission in seasonally dry tropical environment. Environ. Monit. Assess. 158: 197–212.

Sharma, R.K., Agrawal, M.et al. (2008a). Atmospheric deposition of heavy metals (Cu, Zn, Cd and Pb) in Varanasi City, India. Environ. Monit. Assess. 142: 269–278.

Sharma, R.K., Agrawal, M.et al. (2008b). Heavy metal (Cu, Zn, Cd and Pb) contamination of vegetables in urban India: A case study in Varanasi. Environ. Pollut. 154: 254–263.

Sharma, R.K., Agrawal, M.et al. (2009). Heavy metals in vegetables collected from production and market sites of a tropical urban area of India. Food Chem. Toxicol. 47: 583–591.

Shukla, V. and Upreti, D.K. (2007). Effect of metallic pollutants on the physiology of lichen, Pyxine subcinerea Stirton in Garhwal Himalayas. Environ. Monit. Assess. 141: 237–243.

Simmonds, P. G., Derwent, et al. R. G. (2004). Significant growth in surface ozone at Mace Head, Ireland, 1987-2003. Atmospheric Environment 38(28): 4769-4778.

Simpson, D., Benedictow, A. et al. (2012). The EMEP MSC-W chemical transport model – technical description. Atmospheric Chemistry and Physics 12: 7825-7865.

Singh, M., Singh, D.et al. (2008). Aerosol characteristics at Patiala during ICARB-2006. J. Earth Syst. Sci. 117: 407-411.

Singh, P., Singh, S.et al. (2012). Assessment of the interactive effects of ambient O 3 and NPK levels on two tropical mustard varieties (Brassica campestris L.) using open-top chambers. Environ. Monit. Assess. 184: 5863–5874.

Singh, R.P. (2004). Variability of aerosol parameters over Kanpur, northern India. J. Geophys. Res. 109.
 Singh, S., Bhatia, A.et al. (2013). Synergistic action of tropospheric ozone and carbon dioxide on yield and nutritional quality of Indian mustard (Brassica juncea (L.) Czern.). Environ. Monit. Assess. 185: 6517–6529.

Singh, S.P., Satsangi, G.S. et al. (2001). Multiphase measurement of atmospheric ammonia. Chemosphere - Glob. Change Sci. 3: 107– 116.

Sitch, S., Cox, P. M. et al. (2007). Indirect radiative forcing of climate change through ozone effects on the land-carbon sink. Nature 448 (7155): 791-U794.

Slootweg, J., Hettelingh, J.-P. et al. (2010). Critical loads of heavy metals and their exceedances. In: Slootweg, J., Posch, M. et al. (Eds). Progress in the modelling of critical thresholds and dynamic modelling, including impacts on vegetation in Europe. CCE Status Report 2010, Coordination Centre for Effects, RIVM, Bilthoven, The Netherlands.

Somashekaralah, B.V., Padmaja, K.et al. (1992). Phytotoxicity of cadmium ions on germinating seedlings of mung bean (Phaseolus vulgaris): Involvement of lipid peroxides in chlorphyll degradation. Physiol. Plant. 85: 85–89.Smith P.N., Cobb G.P. et al. (2007). Contaminant exposure in terrestrial vertebrates. Environmental Pollution 150: 41-64.

Spranger, T., Hettelingh, J.-P. et al. (2008). Modelling and mapping long-term risks due to reactive nitrogen effects: an overview of the LRTAP Convention activities. Environmental Pollution, 154:482-487.

Steiner, A.L. and Chameides, W.L. (2005). Aerosol-induced thermal effects increase modelled terrestrial photosynthesis and transpiration. Tellus B 57: 404–411.

Steinnes, E., Rühling, Å. et al. (1997). Reference material for large-scale metal deposition surveys. Accreditation and Quality Assurance 2: 243-249.

Stevens, C.J., Duprè, C. et al. (2011). The impact of nitrogen deposition on acid grasslands in the Atlantic region of Europe. Environmental Pollution 159: 2243-2250.

Streets, D.G., Yan, F.et al. (2009). Anthropogenic and natural contributions to regional trends in aerosol optical depth, 1980–2006. J. Geophys. Res. 114.

Sutton, M.A., Howard, C.M. et al. (2011). The European nitrogen assessment: sources, effects and policy perspectives. Cambridge University Press, UK.

Takemura, T. (2005). Simulation of climate response to aerosol direct and indirect effects with aerosol transport-radiation model. J. Geophys. Res. 110.

Tang, H., Takigawa, M.et al. (2013). A projection of ozone-induced wheat production loss in China and India for the years 2000 and 2020 with exposure-based and flux-based approaches. Glob. Change Biol. 19: 2739–2752.

Tiwari, S. and Agrawal, M. (2011). Assessment of the variability in response of radish and brinjal at biochemical and physiological levels under similar ozone exposure conditions. Environ. Monit. Assess. 175: 443–454.

Tripathi, R. and Agrawal, S.B. (2012). Effects of ambient and elevated level of ozone on Brassica campestris L. with special reference to yield and oil quality parameters. Ecotoxicol. Environ. Saf. 85: 1–12.

Tripathi, R. and Agrawal, S.B. (2013). Interactive effect of supplemental ultraviolet B and elevated ozone on seed yield and oil quality of two cultivars of linseed (Linum usitatissimum L.) carried out in open top chambers. J. Sci. Food Agric. 93: 1016–1025.

Twomey, S. (1977). Atmospheric aerosols. Elsevier, Amsterdam, Netherlands.

Vandermeiren, K., Harmens H. et al. (2009) Impacts of ground level ozone on crop production in a changing climate, page 213-243. In: Singh S.N. (Ed.). Climate Change and Crops. Springer, Berlin, pp. 213 – 243. ISBN: 978-3-540-88245-9.

Van Dingenen, R., Dentener, F.J. et al. (2009). The global impact of ozone on agricultural crop yields under current and future air quality legislation. Atmospheric Environment 43: 604–618.

Velissariou, D. and Davison A. W. (1996). The adverse effects of ozone on six Greek wheat varieties and five F1 lines of maize. Phytopathologia Mediterranea. 35: 215.

Velissariou, D., Davison, A.W. et al. (1992). Effects of air pollution on *Pinus halepensis* Mill. 1. Pollution levels in Attica, Greece. Atmospheric Environment 26A: 373-380.

Velissariou, D. and Skretis, L. (1999). Critical levels exceedances and ozone biomonitoring in the Greek fir forest (*Abies cephalonica* loud.) at the Parnis mountain national park in Attica, Greece, page 205- 208. In: Fuhrer, J. and Achermann, B. (Eds.). Critical Levels for Ozone - Level II. UNECE Convention on LRTAP. SAEFL, Bern

Velissariou, D., Venouziou, J. et al. (2000). Cases of necrotic symptoms on Greek crop plants. Is air pollution a cause for many of them? A case study at Gomfi, Trikala, Thessaly. Phytopathologia Mediterranea 39: 322.

Vingarzan, R. (2004). A review of surface ozone background levels and trends. Atmospheric Environment 38(21): 3431-3442. WMO/UNEP, 2011. Integrated Assessment of Black Carbon and Tropospheric Ozone (No. ISBN: 92-807-3141-6). UNEP (DEW/1351/NA).

Working Group on Effects (2013a). Benefits of air pollution control for biodiversity and ecosystem services. 32nd Session WGE, informal document 1.

http://www.unece.org/fileadmin/DAM/env/documents/2013/air/wge/No.1 Benefits of air pollution control for biodiversity and ecosystem_services.pdf

Working Group on Effects (2013b). Guidance document on health and environmental improvements using new knowledge, methodsand data. Document ECE/EB.AIR/2013/8.

http://www.unece.org/fileadmin/DAM/env/documents/2013/air/eb/ECE_EB.AIR_2013_8_E.pdfEW/1351/NA.

Wu, Y.-S., Fang, G.-C.et al. (2007). A review of atmospheric fine particulate matter and its associated trace metal pollutants in Asian countries during the period 1995–2005. J. Hazard. Mater. 143: 511 – 515.

Yamaguchi, M., Otani, Y.et al. (2014). Effects of long-term exposure to ammonium sulfate particles on growth and gas exchange rates of Fagus crenata, Castanopsis sieboldii, Larix kaempferi and Cryptomeria japonica seedlings. Atmosperhic Environ. In press.

Yousefi, N., Chehregani, A.et al. (2009). Effect of the Heavy Metals on the Developmental Stages of Ovule and Seed Proteins in Chenopodium botrys L. (Chenopodiaceae). Biol. Trace Elem. Res. 144: 1142–1149.

Zheng, B., Ma, Y.et al. (2010). Assessment of the influence of global dimming on the photosynthetic production of rice based on threedimensional modeling. Sci. China Earth Sci. 54: 290–297.

Annex 1: Country reports

Here we have included country reports from participants of the programme of the ICP Vegetation. The short reports summarise their research; the responsibility for their content lies with the participants.

Country reports have been submitted by:

- Albania
- Croatia
- Egypt (as outreach to North Africa)
- Greece
- Macedonia
- Romania
- Russian Federation
- Serbia
- Slovenia

Albania

Survey of atmospheric deposition of heavy metals in Albania using mosses

Pranvera Lazo¹, Flora Qarri², Lirim Bekteshi³, Trajce Stafilov⁴, Katerina Baceva⁴ Jani Marka¹

¹Department of Chemistry, Faculty of Natural Sciences, University of Tirana, Albania. <u>pranveralazo@gmail.com</u>

² Department of Chemistry, University of Vlora, Albania.

³Department of Chemistry, Faculty of Natural Sciences, University of Elbasan, Albania.

⁴ Institute of Chemistry, Faculty of Science, Sts. Cyril and Methodius University, Skopje, Macedonia.

Background and aims the study

Albania is a small county (28 000 km²) with a complex topography, climate, geology, and is characterized by high anthropogenic influences. The first study monitoring atmospheric deposition of metals using mosses in Albania was performed within the framework of the ICP Vegetation (Harmens et al., 2010). The main aim of this study is to express the spatial patterns of heavy metals distribution in mosses in 2010/11 in Albania and to identify the main polluted areas in the country.

Materials and methods

The carpet-forming mosses *Hypnum cupressiforme* and *Pseudoscleropodium purum* were collected according to the guidelines set out in the experimental protocol for the 2010/11 survey (ICP Vegetation, 2010). The distribution of the sampling sites throughout Albania was shown in a previous publication (Qarri et al., 2013). Ten elements (AI, As, Cd, Cr, Cu, Fe, Ni, Pb, V and Zn) were measured in moss samples collected from 62 sampling sites across Albania during the dry autumn and summer period of 2010 and 2011. The total digestion of moss samples was done according to the method presented by Barandovski et al. (2008). The quality of the data was checked by multiple analyses of samples and by analyzing the certified moss reference materials M2 and M3 (Steinnes et al., 1997; Harmens et al., 2010).

Results and Discussions

The 2010 data on the concentration of 10 elements in 62 moss samples from Albania are summarized below, using descriptive statistics (**Table 1**) and multivariate analysis (Pearson Correlation, **Table 2**).

Parameters	Fe	Ni	V	Al	Cr	As	Pb	Cd	Cu	Zn
Range	469–5488	1.6-131	1.15-16.9	535-6974	1.6-31.7	0.05-2.9	1.3–19.7	0.038-0.89	2.14-16	1.0–68
Mean	1892	11.36	4.23	1958	6.38	0.541	3.28	0.17	6.07	14.06
Median	1618	5.85	3.51	1638	4.75	0.305	2.41	0.107	5.58	13.8
St dev.	1105	19.3	2.79	1178	5.39	0.64	3.21	0.16	2.8	11.6

Table 1Descriptive statistics of the element concentrations (mg/kg, DW) in mosses sampled (n=62) in
Albania in 2010.

The median values of Cr, Fe, Ni, V, Zn and Al are similar to those of neighbouring countries (Qarri et al. 2013), but higher than those of European countries (Harmens et.al. 2013). The highest values of these elements were measured near industrial centres (the central part of the country). To distinguish between lithogenic and anthropogenic origin of the elements in mosses, correlation analysis was carried out (Table 2). Good correlations (R²>0.5, P>0.001) were found between Fe and Cr, Ni, V, Al, which can be explained by their lithogenic (Fe-Al, V correlation) and geogenic (Fe-Cr, Ni correlation) origin.

Aluminium, chromium, iron, nickel and vanadium

The background level of Al in Albanian moss samples was higher than in other European countries (Harmens et al., 2013). The highest Al concentration was found in the south and in central part of Albania. The main contribution of Cr, Fe, Ni and V elements is coming from the Elbasan ferrochromium metallurgical plant (Lazo et al., 2013) and the mining industry in Albania. A high level of wind-blown dust occurs in the south and a high level (for most metals) of industrial activity is present in the mid-east of Albania. The association of Cr and Fe is also related to current air pollution (Lazo et al. 2013). Their highest concentration is present near the ferrochromium metallurgy in Elbasan town and chromites deposition areas of Albania.

	Cd	As	Cr	Cu	Pb	Ni	V	Zn	Al	Fe
Cd	1									
As	0.19	1								
Cr	0.46^{1}	0.18	1							
Cu	0.39^{2}	-0.05	0.31	1						
Pb	0.35^{3}	0.08	0.26	0.45^{1}	1					
Ni	0.21	-0.01	0.64 ¹	0.14	0.06	1				
V	0.30	0.34^{3}	0.51 ¹	0.30	0.31	0.13	1			
Zn	0.43 ¹	-0.09	0.43	0.62 ¹	0.52 ¹	0.21	0.34^{3}	1		
Al	0.34 ³	0.60 ¹	0.43 ¹	0.11	0.12	0.08	0.67 ¹	0.02	1	
Fe	0.43 ²	0.44^{1}	0.81 ¹	0.38 ²	0.30	0.51 ¹	0.72 ¹	0.39	0.75 ¹	1

 Table 2
 Pearson Correlation Coefficients between element concentrations in mosses in Albania in 2010.

¹ P<0.001, ² P<0.005, ³ P<0.01; bold – Pearson Correlation Coefficient > 0.50.

Arsenic, cadmium, copper and zinc

Arsenic concentrations in mosses were generally low in the western part and higher in eastern part of Albania, but lower than in neighbouring countries (Harmens et al., 2013). Cu, Cd and Zn concentrations were generally low in mosses sampled in Albania compared to many other European countries (Harmens et al., 2013). Road transport may have a considerable effect on the distribution of these elements in air pollution in Albania.

Conclusions

Moss biomonitoring provides a cheap, complementary method to deposition analysis for the identification of areas at risk from high atmospheric deposition fluxes of heavy metals. Based on the median distribution of Albanian data of moss survey 2010/2011 we suggest that the elements Zn, Cd, Cu, Pb and As do not reach high enough concentrations as a result of long-range atmospheric transport and deposition to cause adverse effects on terrestrial ecosystems. However, the elements AI, Fe, Cr, Ni and V appear to have the highest median values among European countries and this may result in effects on vegetation due to the accumulative nature of heavy metals in soils and vegetation. Soil dust, industry emissions, waste incineration and road traffic were identified as main factors causing air pollution from heavy metals in Albania.

References

- Barandovski, L., Cekova, M., Frontasyeva, M.V., Pavlov, S.S., Stafilov, T., Steinnes, E., Urumov, V. (2008). Atmospheric deposition of trace element pollutants in Macedonia studied by the moss biomonitoring technique. Environmental Monitoring and Assessment 138: 107-118.
- Harmens H, Norris Ď, Mills G. and the participants of the moss survey (2013). Heavy metals and nitrogen in mosses: spatial patterns in 2010/2011 and long-term temporal trends in Europe. ICP Vegetation Programme Coordination Centre, Centre for Ecology and Hydrology, Bangor, UK, 63 p. <u>http://icpvegetation.ceh.ac.uk</u>
- Harmens, H., Norris, D.A., Steinnes, E., Kubin, E., et al. (2010). Mosses as biomonitors of atmospheric heavy metal deposition: spatial and temporal trends in Europe. Environmental Pollution 158: 3144-3156.
- ICP Vegetation (2010). Heavy Metals in European Mosses: Survey. Monitoring Manual. International Cooperative Programme on Effects of Air Pollution on Natural Vegetation and Crops, <u>http://icpvegetation.ceh.ac.uk/manuals/ documents/ UNECEHEAVYMETALSMOSSMANUAL2010POPsadaptedfinal_220510_.pdf</u>
- Lazo, P., Bekteshi, L., Shehu, A. (2013). Active moss biomonitoring technique for atmospheric deposition of heavy metals study in Elbasan city, Albania, Fresenius Environmental Bulletin 23 No 1a: 213-219.
- Qarri, F., Lazo, P., Stafilov, T., Frontasyeva, M., Harmens, H., Bekteshi, L., Baceva, K., Goryainova, Z. (2013). Multi-elements atmospheric deposition study in Albania. Environmental Science and Pollution Research, October 2013.
- Steinnes, E., Rühling, Å., Lippo, H., Makinen, A. (1997). Reference materials for large-scale metal deposition surveys. Accreditation and Quality Assurance 2: 243-249.

Croatia

- Lead institute: Zdravko Spiric and Vladimir Kusan, Oikon Ltd. Institute for Applied Ecology, Trg senjskih uskoka 1, Zagreb, Croatia. zspiric@oikon.hr
- **Collaborating institutes:** Trajce Stafilov and Ivana Vuckovic, Institute of Chemistry, Faculty of Science, Ss. Cyril and Methodius University, Skopje, Macedonia. <u>trajcest@pmf.ukim.mk</u>
 - Mariana Frontasyeva, Frank Laboratory of Neutron Physics, Joint Institute for Nuclear Research, Dubna, Russian Federation. <u>marina@nf.jinr.ru</u>
 - Delko Barisic and Branko Vekic, Ruđer Bošković Institute Zagreb, Croatia
 - Marin Glad, Teaching Institute of Public Health, Primorsko-Goranska County, Rijeka, Croatia. marin.glad@zzjzpgz.hr

Background

Since 2006, Croatia has participated in the European moss survey. Moss samples were collected in 2006 and 2010 on a nearly regular grid of 23 km x 23 km. Moss samples were collected during the summer of 2006 and summer/autumn of 2010 (**Figure 1**), from 98 and 121 locations respectively, evenly distributed over the country with additional samples taken in/around urban/industrial areas. The most dominant moss species were *Hypnum cupressiforme, Pleurozium schreberi, Brachythecium rutabulum* and *Homalothecium sericeum*. This study was undertaken in order to provide an assessment of air quality throughout Croatia and to generate information needed for better identification of local pollution sources as well as transboundary pollution, and improving the potential for assessing environmental and health risks in Croatia associated with dry and wet deposition of toxic metals.





Metals, nitrogen & radionuclide concentrations in mosses

Here we present the results obtained in the 2010 moss survey in Croatia and compare these results with those obtained in the previous survey in 2006, in order to evaluate spatial patterns and temporal deposition trends. As an example, the spatial pattern of Pb concentrations in mosses in 2010 is shown (**Figure 2**). The content of 21 elements was determined by ICP-AES and atomic absorption spectrometry (AAS). Principal component analysis (PCA) was applied in order to show association between the elements. Six factors (F1 to F6) were determined, of which two are anthropogenic (F3 and F6), two are mixed geogenic-anthropogenic (F1 and F5) and two are geogenic factors (F2 and F4). In addition, 22 out of 121 representative moss samples were subjected to gamma-spectrometric analyses for assessing the activity of the naturally occurring radionuclides (data not shown). In 2010, the nitrogen concentration in mosses was determined for the first time using the Kjeldahl method (Figure 4). From data obtained in 2010 (Špirić et al., 2013), it can be concluded that the median values and ranges of all elements obtained in this study are very similar to the median values and ranges obtained in the previous study in 2006 (Špirić et al., 2012). Only a few elements (Cd, Cu, Mg, Ni and Pb) have a slightly higher median values were recorded in 2010.

Table 1

Comparison of the results of the 2006 and 2010 Croatian moss survey.

	Cro	atia 2010	Croa	atia 2006	
No. of samples	r	121 =		n= 94	
Element	Median	Range	Median	Range	
Ag	0.032	0.001-0.155	-	-	
Al	878	112-4493	1350	398-2146	
As	0.36	0.05-1.00	0.37	0.10-6	
Ва	20.64	4.49-94.30	32	7-192	
Ca	6632	2649-20795	7623	2832-26740	
Cd	0.38	0.10-1.42	0.27	0.07-1.9	
Cr	1.94	0.41-8.55	2.8	0.76-33	
Cu	8.53	4.72-22.69	7.5	3.7-22.7	
Fe	789	85.00-4028	1000	320-12140	
Hg	0.043	0.010-0.145	0.064	0.007-0.301	
К	3891	1552-9279	8085	2565-23720	
Li	0.55	0.11-4.27	-	-	
Mg	3059	1619-4740	2120	676-12740	
Mn	99.1	16.10-928	106	20-1421	
Na	120	65.00-304	169	67-2332	
Ni	3.16	1.04-14.66	2.7	0.66-18	
Р	1134	419-3117	-	-	
Pb	3.21	1.11-36.64	2.46	0.06-82.4	
Sr	16.00	4.74-54.03	21	4-125	
V	2.55	0.23-37.26	3.1	0.91-32	
Zn	24 80	11 64-77 13	29	12-283	



Figure 2 Distribution of Pb (left) and N concentrations (right) in mosses in 2010.

Discussion and conclusion

From the moss data it can be concluded that anthropogenic heavy metal pollution has not changed significantly in Croatia between 2006 and 2010. The main anthropogenic sources of heavy metals are light and heavy industry, transportation, steel industry, textile industry, thermoelectric plant and oil deposits and refineries whose activities are carried out near big industrialized cities such as Zagreb, Sisak, Rijeka, Kutina, Split and Sibenik. Moss samples collected near these regions showed the highest content of typical air pollution from heavy metals. In comparison with the results obtained in other European countries, it can be concluded that Croatia is more polluted than many parts of northern and western Europe, but less polluted than many of its neighbouring countries. There is a clear need to continue the moss survey in order to assess air quality throughout Croatia and to produce information needed for better identification of pollution sources and assessment of environmental and health risks associated with dry and wet deposition of toxic metals in Croatia.

- Špiric Z, Frontasyeva M, Steinnes E & Stafilov T (2012). Multi-element atmospheric deposition study in Croatia, International Journal of Environmental Analytical Chemistry 92: 1200-1214.
- Špiric, Z., Vuckovic, I., Stafilov, T., Kusan, V., & Frontasyeva, M. (2013). Air Pollution Study in Croatia Using Moss Biomonitoring and ICP-AES and AAS Analytical Techniques. Archives of Environmental Contamination and Toxicology 65: 33-46.

Egypt*

Samia Madkour, Damanhour University, Damanhour

Lead institute: 1) The Egyptian Environmental Affairs Agency (EEAA), Ministry of State for Environmental Affairs (MSEA), Cairo, Egypt. www.eeaa.gov.eg [Ahmed Abou Elseoud, Air Quality Dept.].

2) National Research Center, El-Dokki, Cairo,Egypt. <u>info@nrc.sci.eg</u>. [Nasser M. Abdellatif, Air Pollution Department (Ozone + Heavy metals)]

Collaborating Institute: Institute of Environmental Studies and Research, Ein-Shams University, Abbassia, Cairo [Ibrahim M. El-Gamal (Heavy metals only)]

Background

Air pollution is one of the most important challenges and obstacles facing development in Egypt at present. Air pollution emissions pose an increasing threat to agriculture. High rates of consumption of fossil fuel caused by the extensive urbanization, industrial development and the increase in motorized vehicular traffic coupled with the rapid growth in population during the last few decades have enhanced the overall air pollution level.

As the control of air pollution represents one of the primary concerns of the Ministry of State for Environmental Affairs (MSEA) and the Egyptian Environmental Affairs Agency (EEAA), continuous efforts are made for enforcing existing environmental legislation and dealing with the air quality problem (Law 4/1994 for the Environment and its amendment law 9/2009). Accordingly, a comprehensive national air quality monitoring system has been established as part of the Environmental Information and Monitoring Program (EIMP) of EEAA as a long-term commitment to this issue. The monitoring network started with 38 stations and was initially implemented with support from the Danish Government (MSEA/EEAA Air Quality Report, 2001). The monitoring system has been operational between the years 1998-2009, measuring concentrations of common air pollution parameters such as sulphur dioxide (SO₂), nitrogen dioxide (NO₂), carbon monoxide (CO), ozone (O₃), particulate matter (PM10) and lead (Pb). This effort was carried out by monitoring stations distributed throughout the country. In 2001, EEAA took over full responsibility for the operation and maintenance of the system and the number of stations increased gradually to a total of 87 monitoring sites in 2007. This report will focus on O₃, NO₂ and Pb which have been known to have a great impact on agriculture and therefore have been the most studied. Despite the differences between the monitoring locations, levels of the ambient gaseous pollutants O_3 (25-92 ppb) and NO_2 (32-62 ppb) are high enough to have significant impacts on the growth and yield of local varieties of crops (Egypt State of the Environment Reports, 2008 & 2009).

Egyptian agriculture is characterized by the limited cultivable area, mainly along the Nile banks and the Delta since the river is almost the single main source of irrigation water. The cultivated area represents 3.7% of the total area of Egypt (1 million km²). It is considered as one of the most intensive agricultural systems in the world. The agricultural surface unit is cultivated two or three times a year, thus, intensifying the harvested area 2-3 folds above the cultivated area. Economically important crops which are at risk from air pollution include cereals (e.g. wheat, maize & rice), fodder (clover & alfalfa), legumes (beans & lentil), fibre (cotton) and vegetables (e.g. tomato, potato & onion).

Main air pollution problems

Ozone. The increases in air pollution that have occurred around the urban industrial centres of Cairo and Alexandria in Egypt are particularly problematical since these are in the same location as the primary agricultural region, which is limited to the Nile river basin and the Delta. Studies of the effects of O_3 pollution on vegetation have been carried out in the last 20 years in the greater Cairo area and around the main roads within the Nile Delta and Upper Egypt region. Temporal and spatial patterns were followed and recorded by the MSEA/EEAA. Results of these efforts are presented in **Figure 1**. Ozone levels recorded were consistently greater than the universally accepted 40ppb vegetation damage threshold. Hourly means of O_3 further exceeded the above levels and reached 100 ppb or more depending on the site, the weather conditions and the season. Visible injuries included necrosis, black or red spots and chlorosis especially on bioindicator species such as Jew's mallow, white clover (60 % and 54% leaves injured respectively), lettuce and rocket (Madkour & Laurence, 2002).

^{*} As outreach to North Africa



Figure 1 Temporal (left) and spatial (right) trends of the annual average ozone concentrations in Egypt.

Controlled environmental studies aiming to determine the sensitivity of local Egyptian crops and cultivars to O_3 were conducted since 1994. Sensitive species included radish and turnip (Hassan et al., 1995), common bean (Madkour, 1998 & Madkour et al., 2011), clover (El-Shamy and Madkour, 2008) and tomato (Madkour and Abou Salem, 2014).

The impact of O_3 on the growth and yield of local varieties has been assessed. Plant biomass of radish (Raphanus sativus L. cv. Balady) and turnip (Brassica rapa L. cv. Sultani) were reduced at suburban and rural sites in Alexandria. The study proved that levels of ambient O_3 in Egypt are high enough to have significant impacts on the growth and yield of local varieties of vegetable crops, even at a time of the year when O₃ levels are relatively low (Hassan et al., 1995). Open Top Chambers (OTC) experiments were conducted to study O₃ effects on the yield of an Egyptian cultivar of wheat. Results showed that exposure to 61 ppb O₃ caused 60% decrease in the total grain dry weight/plant and 20-48% decrease in other yield parameters (Hassan et al., 1999). O₃-induced yield reductions have also been observed with other economically important crops. Estimated yield losses due to ozone exposure in the range 77-166 ppb at four different open field sites were proportional to the ozone level at each site. Reductions in yield of wheat (Triticum aestivum L. cv. Giza 68), broad bean (Vicia faba L. cv. Lara), kidney bean (Phaseoulus vulgaris L. cv. Giza 3) and pea (Pisum sativum L. Perfection) were 9-46%; 13-33%; 20-45%; 3-30% respectively (Ali et al., 2008). The combined impact of O_3 , NO_2 and SO_2 in ambient air were assessed using three cultivars of pea at urban (90 ppb O_3 , 24 ppb NO₂ & 30 ppb SO₂) and rural (76 ppb O₃, 12 ppb NO₂ & 13 ppb SO₂) sites in Sharkia Province north east of Egypt. Losses in growth and decrease in yield of pea plants reached 40% and 10% respectively, at the urban site and were less at rural sites (Ali, 2004).

Nitrogen dioxide. Results of the air monitoring network (1999-2990, **Figure 2**) show that recorded NO₂ concentrations in ambient air has always exceeded the maximum allowable standard set by the World Health Organization (40 μ g/m₃). It is noteworthy that this problem is not a new one, the increase in vehicle number in recent years led to a rise in the average annual concentrations of NO₂. In addition, the expansion in using natural gas either in industry, production of electricity or as fuel for vehicles contributed in increasing NO₂ concentrations.



Figure 2 Temporal (left) and spatial (right) trends of the annual average concentrations of nitrogen dioxide in Egypt.

Lead. Monitoring results for lead (Pb) concentrations in Greater Cairo (1999- 2009) show a significant decrease (0.4-0.7 μ g/m³) during 2007 and 2008 when compared to concentrations observed between 2000 and 2002 (1.7 μ g/m³; **Figure 3**). This reduction in Pb was a result of the exerted efforts of the MSEA to implement a National Programme aimed at reducing Pb pollution loads in Shoubra El-Khaimah, an extensive industrial area within the greater Cairo region. This programme began in 1998 and ended in March 2008 and consisted of a project to transfer all the foundries and cleaning their

lead-contaminated sites in Shoubra al-Khaimah; the expansion in producing unleaded gasoline by the Ministry of Petroleum; and the gradual replacement of gasoline with compressed natural gas as fuel for public transport. These efforts had been accompanied with MSEA amendment of Law 4 /1994 with respect to the permissible limits of lead concentrations in air to become 0.5 μ g/m³ in residential areas and 1.5 μ g/m³ in industrial areas instead of the general limit of 1 μ g/m³, in both industrial and residential areas. Bio-accumulator plants (*Lolium multiflorum* L.) were used to assess Pb and Cd levels in plants in great Cairo. Evidence presented show that both Pb and Cd levels decreased with distance from highways, traffic volume and the existence of green barriers (El-Gamal, 2000).





Lead concentrations collected on leaf surfaces were high near industrial sites and lower at urban and suburban locations (Abou –El Saadat et al., 2011). Efforts to measure the accumulation of Pb in soils showed that at El-Fayoum Governorate the concentration was below the maximum permissible limits and were highest in soil surface especially near the main roads (Abdel-gawad et al., 2007).

Conclusions

- Air pollutant emissions pose an increasing threat to agriculture in Egypt;
- There is a great need to assess the current and future impact of air pollution and to develop air quality guidelines for agriculture in Egypt, related to the local cropping pattern and climate;
- Air pollution monitoring has tended to focus on urban areas, and there is a requirement for more rural monitoring on a regular basis.

References

- Abd Elgawad, M., Mahmoud, H.A.A., Shendi, M. and Ghabour, S.I. (2007). Status of some heavy metals in Fayoum district soils. Proceeding of the 3rd Conference for Sustainable Agricultural Development, Facity of Agriculture, EI-Fayoum University (12-14 Nov.) pp. 507-526.
- Abou El Saadat, E.M., Hassan, M.R., Hassan, I.A.and Weheda, B.M. (2001). Heavy metal content in leaves of Fiscus retusa collected from contaminated and uncontaminated sites in Northern Egypt: Mitigation of toxicity by washing treatments. Universal Journal of Env. Research and Tech. 1(4): 408-415.
- Ali, A. A. (2004). An open top chambersstudy to evaluate the impact of airborne pollutants on the growth of pea (*Pisum sativum* L.) cultivars at three sites of Sharkia Province, Egypt. Egyptian Journal of Biology 6: 52-61.
- Ali, A., Alfarhan, A., Robinson, E., Bokhari, N., Al-Rasheid, K. and Al-Koraishy, S. (2008). Tropospheric ozone effects on the productivity of some crops in central Saudia Arabia. American Journal of Environmental Sciences 4(6): 631-637.
- El-Gamal, I.M. (2000). Distribution pattern of some heavy metals in soil and plants along El-Moukattam highway. ICEHM, Cairo University, Egypt.pp. 518-524.
- El-Shamy, Sawsan and Madkour, Samia (2008). Some physiological and biochemical changes in economic crops caused by the air pollutant ozone. M SC. Thesis. Alexandria University, Alexandria, Egypt.
- Hassan, I., Anttoneen, S., Ashmore, M., Bell, J.N.B., Bender, J. & Weigel, H.J. (1999). Effect of ozone on ethylene biosynthesis and yield of wheat (*Triticum aestivum* L.). Pak. J. Biol. Sci. 332 335
- Hassan, I.A., Áshmore, M.R. & Bell, J.N.B. (1995). Effects of O₃ on radish and turnip under Egyptian field conditions. Environ Pollut. 89: 99 108.
- Madkour, S.A. and Laurence J.A. (2002). Egyptian plant species as new ozone indicators. Environmental Pollution 120:339-353.
- Madkour, S.A., Abou-Salem, A. (2008). The role of salicylic acid and jasmonate in conferring tolerance against ozone injury in tomato plants. (In preparation).
 Madkour, S.A., Abou-Salem, A. (2014). The role of salicylic acid and jasmonate in conferring tolerance against ozone injury in
- Madkour, S.A., Abou-Salem, A. (2014). The role of salicylic acid and jasmonate in conferring tolerance against ozone injury in tomato plants. Proceedings of the 27th Task Force Meeting of the UNECE ICP Vegetation, Jan. 28 30, 2014 Paris, France.
- Madkour, S.A., Saitanis, K., Saad, E.M. (2011). Egyptian bean genotypes as ozone bioindicators. Proceedings of the 24th Task Force Meeting of the UNECE ICP Vegetation, Jan. 31st Feb.2nd,2011 Rapperswil, Switzerland.
 Madkour, Samia A. (1998). Defense mechanisms against ozone injury in beans (*Phaseolus vulgaris* L.) genotypes differing in
- Madkour, Samia A. (1998). Defense mechanisms against ozone injury in beans (*Phaseolus vulgaris* L.) genotypes differing in ozone tolerance. J. Agric. Sci. Mansoura Univ. 23(7): 3203-3225.
- MSEA/EEAA (2008). Egypt State of the Environment Report.
- MSEA/EEAA (2009). Egypt State of the Environment Report.

Greece

Phytotoxicity of tropospheric ozone

Institutes: Dimitris Velissariou, Technological Educational Institute of Peloponnese (1999 - today), Benaki Phytopathological Institute (1988-1999). <u>d.velissariou@teikal.gr</u> Costas J. Saitanis, Karandinos M.G., Riga-Karandinos A.N., Lekkas D., Agathokleous E., Agricultural University of Athens, Laboratory of Ecology and Environmental Sciences, Iera Odos 75, Votanikos, Athens, Greece. <u>saitanis@aua.gr</u>

Background

Tropospheric ozone is the major air pollutant in major Greek cities (Athens, Thessaloniki, Patra, Volos, etc.) these days. There is also evidence that ambient ozone occurs at increased levels in many rural places across the country. This happens because Greece, along with other Mediterranean countries, is characterized by long periods of strong sunlight favouring photochemical pollution formation. Air pollution effects on plants have been studied in Greece since the mid seventies, mainly from point source air pollutants (airborne fluoride) (Holevas & Velissariou, 1986). In 1988, the first ozone Bel-W3 tobacco bio-monitoring campaign was carried out in Greece, at 23 sites in Attica, as well as a visible injury survey on Aleppo pine (*Pinus halepensis* Mill.) at 18 sites and short term ozone monitoring in two rural areas (Velissariou et al., 1989, 1990, 1992, Velissariou, 1993). In addition, an ozone exposure study was conducted on main Greek crops in fumigation chambers at Newcastle University (U.K.) from 1988 to 1990 (Velissariou, 1993, Velissariou & Davison, 1996). In recent years, tropospheric ozone impacts have been studied in major Greek cities (Saitanis et al., 2004; Riga-Karandinos & Saitanis. 2005; Riga-Karandinos et al., 2006), as well as in many rural places across the country (Saitanis & Karandinos, 2001; Saitanis, 2003, 2008; Kalabokas & Reparis, 2004).

Impacts of ozone on vegetation

Severe damages on agricultural crops (yield losses and/or commercial value losses) due to ozone toxicity have been reported in Greece since 1993, on crops like wheat, maize, cotton, potato, watermelon, muskmelon, beans, lettuce (**Figure 1**), onion, fodder crops (Velissariou, 1996, 1999; Velissariou et al., 2000; Fumagali et al., 2001). These damages usually occur during short term acute photochemical episodes (Figure 1), common in Mediterranean climates, all over the year and particularly in areas favouring temperature inversions. Transboundary ozone is also being detected in Greece, probably affecting crop plants and natural flora in the long term. Moreover, ozone toxicity symptoms have been detected on the needles of Greek fir (*Abies cephalonica* loud.) and high ozone concentrations have been measured in two Greek fir forest ecosystems: Parnis mountain in Attica (Velissariou & Skrekis, 1999) and Taygetos mountain in Peloponnese southern Greece, (Velissariou & Salmas, 2008). Bioindicator campaigns with Bel-W3 tobacco and white clover (**Figure 2**) have also shown high ozone levels in other mountainous regions (Central Peloponnese and Pelion Mountain, Thessaly) (Saitanis et al., 2004, 2006), causing visible leaf injury.











Discussion, conclusions, recommendations

Photochemicals and particularly tropospheric ozone are the dominating air pollution problem for vegetation in Greece in the last decades. High levels of tropospheric ozone due to the climatic conditions in Mediterranean countries, including Greece, such as strong and prolonged solar irradiation, favour ozone formation. Complex topography (valleys and basins) causing temperature inversions enhance the problem. Irrigated crops (vegetables, arable crops, etc.) are at high risk of ozone exposure due to high ozone uptake when water supply is non-limiting. Severe damages on agricultural crops (yield losses and/or commercial value losses) have been reported for the most important agricultural areas in the country (Thessaly, Peloponnese, Crete, etc.). Moreover, there is evidence that ozone may play important role to the decline of sensitive Greek fir (Abies cephalonica loud.) forest ecosystems.

References

Fumagalli, I., Gimeno, B. S., Velissariou, D., De Temmerman, L., Mills, G. (2001). Atmospheric Environment 35: 2583-2587. Holevas C.D. and Velissariou D.E. (1986). Airborne Fluoride injury to Vegetation in Greece. In : "How are effects of air pollutants on agricultural crops influenced by interaction with other limiting factors". CEC, Concerted Action 'Effects of Air Pollution on Terrestrial and Aquatic Ecosystems', pp 119-123, RISO, Denmark, 1986.

Kalabokas and Reparis (2004). Atmospheric Chemistry and Physics 4:1139–1147.

- Riga-Karandinos and Saitanis (2005). Chemosphere 59:1125-1136.
- Riga-Karandinos et al (2006). International Journal of Environment and Pollution 27(4): 300-312.

Saitanis (2003). Chemosphere 51(9):913-923.

Saitanis (2008). Italian Journal of Agronomy 3(1):71-77.

Saitanis and Karandinos (2001). Chemosphere 44:813-821.

- Saitanis et al. 2004. Bulletin of Environmental Contamination and Toxicology 72(6):1268-1277.
- Velissariou D. (1996). Phytopathol. Mediterr. 35: 228.
- Velissariou D. (1999). In : Critical Levels for Ozone Level II. UNECE Convention on LRTAP, Edited by : Fuhrer J. & Acherman B. Published by : SAEFL, Bern, 1999 (UM-115-E). pp 253-256.

Velissariou D. and Davison A. W. (1996). Phytopathol. Mediterr. 35: 215.

- Velissariou D. and Salmas I. (2008). Phytopathol. Mediterr. 47: 162.
- Velissariou D. and Skretis L. (1999). In : Critical Levels for Ozone Level II. UNECE Convention on LRTAP, Edited by : Fuhrer J. & Acherman B. Published by : SAEFL, Bern, 1999 (UM-115-E). pp 205-208.
- Velissariou D., Barnes, J., Davison A., et al. (1990). Effects of ozone on crops and forests in Attica. In : Environment : air pollution and its effects on vegetation water and soil. Spanish-British Workshop. Madrid 2-5 Oct. 1990.

Velissariou D., Davison A.W., Barnes J.D., et al. (1992). Atmospheric Environment 26A: 373-380..

Velissariou D., Venouziou J. and Papas K. (2000). Phytopathol. Mediterr. 39: 322.

Velissariou D.E. (1993). Ozone sensitivity of important crop plants around Athens, Greece. PhD Thesis, University of Newcastle Upon Tyne, UK.

Velissariou D.E., Davison A. and Ollerenshaw J. (1989). Ozone in Greece. Summaries of Reports of the Committee for Air Pollutions Effects Research (CAPER), Manchester University, April 1989.

Macedonia

- Lead institute: Trajče Stafilov (Coordinator), Lambe Barandovski, Katerina Bačeva, Faculty of Natural Sciences and Mathematics, Ss. Cyril and Methodius University, POB 162, 1000 Skopje, Macedonia. <u>trajcest@pmf.ukim.mk; lambe@pmf.ukim.mk; k.baceva@gmail.com</u>
- **Collaborating institutes:** Marina V. Frontasyeva, Frank Laboratory of Neutron Physics, Joint Institute for Nuclear Research, str. Joliot-Curie, 6, 141980 Dubna, Moscow Region, Russian Federation. <u>marina@nf.jnr.ru</u> Robert Šajn, Geological Survey of Slovenia, Dimičeva ul. 14, 1000 Ljubljana, Slovenia. Robert.Sajn@geo-zs.si

Background

The Republic of Macedonia was involved in the European moss survey under UNECE ICP Vegetation for the first time in 2002, when the atmospheric deposition of trace elements was studied in the whole country. The moss survey was also performed again in 2005 and 2010 using samples of terrestrial mosses *Hypnum cupressiforme* Hedw. and *Homalothecium sericeum* (Hedw.) B.S. & G. Moss samples were collected at 73 sites in 2002 and at 72 sites in 2005 and 2010 evenly distributed over the area of the country, using a dense net of 17x17 km² in accordance with the sampling strategy of the European moss survey programme. The analyses of 42 elements were performed by using of neutron activation analysis (NAA) and atomic absorption spectrometry (AAS) as analytical techniques. The most important emission sources were determined (smelters and drainage systems near the towns of Veles, Tetovo, Kavadarci and Radoviš, as well as mines Sasa, Toranica and Zletovo in the east), and some uranium deposition patterns were identified from the activity of power plants (near Bitola and Kičevo) using lignite coal as fuel.

Concentrations of heavy metals and nitrogen

To determine the content of various elements in the mosses collected in 2010, AAS, ICP-AES and NAA were used. The content of 18 elements (AI, Ba, Ca, Cd, Co, Cr, Cu, Fe, K, Li, Mg, Mn, Ni, P, Pb, Sr, V and Zn) was determined by using AAS and ICP-AES. The analyses were performed at the Institute of Chemistry, Faculty of Natural Sciences and Mathematics, Skopje, Macedonia, while NAA was performed at the Joint Institute for Nuclear Research, Frank Laboratory of Neutron Physics, Department of Neutron Activation Analysis in Dubna, Russian Federation. The content of 42 elements was determined: (Na, Mg, Al, Cl, K, Ca, Sc, Ti, V, Cr, Mn, Fe, Co, Ni, Cu, Zn, As, Se, Br, Rb, Sr, Zr, Mo, Ag, In, Sb, I, Cs, Ba, La, Ce, Sm, Eu, Tb, Dy, Yb, Hf, Ta, W, Au, Th, and U). For the determination of the content of N, the Kjeldahl analytical method was applied.

Results and discussion

Analysing the trend of the median values calculated for the three moss surveys in Macedonia, it can be seen that in the last survey lower median values are observed for all elements, usually related to heavy metal pollution. The analysis with AAS as well as ICP-AES (2010 survey) is based on nitric acid digestion, and in non-destructive NAA determination (2002 and 2005 surveys) the total amount of the elements in question is analysed. The lower median values for Al, Ba, Ca, Cr, Cu, Fe, K, Mg, Mn, Na, Ni, Sr, V and Zn in 2010 can be explained by the difference of the analytical techniques and potentially not determining the total amount of elements using nitric acid digestion, due to their refractory minerals. Due to the intensive ferronickel production in Kavadarci region, the median value for nickel, in the 2005 survey is 2.5 fold higher than the data obtained for 2002. There are cases when the content of some anthropogenic elements are lower in the 2005 survey than those reported for 2002 (As, Cr, Cu, Sb, Se). For example, the newly introduced protective measures on the slag dump from the closed ferrochromium smelter located near Tetovo, contributed a decrease in the content of Cr in the samples from 2005 compared to 2002.

To explain the variation and to reveal associations of chemical elements all moss samples were examined by multivariate analysis. Five factors were identified and 80.3% of the variability of the investigated elements is explained. Factors were identified by visual inspection of similarities of spatial distribution of element patterns, the correlation coefficient, comparison of basic statistical parameters and the results of multivariate analyses. **Factor 1** (As, Al, Ce, Cs, Dy, Fe, Hf, La, Li, Mg, Na, Nd, Yb, Sc, Sm, Ta, Tb, Th, Ti, U, V, W, Zr) represents chemical elements that are naturally distributed near to the region of Bitola, which is affected by the fly ash from the power plant using coal as a fuel and where the content of U and Th is higher than in the other regions. The content of these elements in mosses is significantly influenced by mineral particles released into the atmosphere by wind erosion of local sources or particles attached to the moss in the periods when the soil surface is

covered by water. Factor 2 (Ni, Cr, Co) represents a mixed (geogenic-anthropogenic) association of elements. These elements are affected primarily by natural factors such as lithological background, but are also affected by anthropogenic influence (ferronickel smelter plant near the town of Kavadarci). Factor 3 (Cd, Pb, Sb, Zn) represents the second anthropogenic geochemical association of elements due to the activities of three Pb-Zn mines in the eastern part of the country and due to the pollution from the former Pb-Zn smelter plant in the town of Veles. Factor 4 (Sr, Ba) represents another geogenic association of elements with the highest factor scores in the north-eastern part of the country. Factor 5 (I, Br) indicates marine influences and the highest values of these elements are present in the south-central and eastern part of the country. The spatial distribution of Factors scores is shown in Figure 1.





Spatial distribution of Factor 1 (As, Al, Ce, Cs, Dy, Fe, Hf, La, Li, Mg, Na, Nd, Yb, Sc, Sm, Ta, Tb, Th, Ti, U, V, W, Zr), Factor 2 (Ni, Cr, Co), Factor 3 (Cd, Pb, Sb, Zn), Factor 4 (Sr, Ba) and Factor 5 (I, Br) scores for element concentrations in mosses.

Nitrogen was also analyzed in the moss samples collected in 2005 and 2010 survey. The median value of nitrogen in moss samples was 1.21% (range: 0.70 - 1.54%) in 2005 and 1.06% (range: 0.68 - 1.75%) in 2010.

The results obtained from the investigation show that in comparison with similar studies made in neighbouring countries the pollution situation in the Republic of Macedonia is more favourable, but comparison with more pristine territories in other parts of Europe still shows that the country is exposed to considerable atmospheric metal pollution. Trends of lower median content in the mosses is noticeable in recent years for some elements, due to closing of some of the smelters, using non-leaded fuels and some protective measures taken for some of the dump slugs in the country.

References

- L. Barandovski, M. Cekova, M. V. Frontasyeva, S. S. Pavlov, T. Stafilov, E. Steinnes, V. Urumov (2008). Atmospheric deposition of trace element pollutants in Macedonia studied by the moss biomonitoring technique, Environmental Monitoring and Assessment 138: 107-118.
- L. Barandovski, M. V. Frontasyeva, T. Stafilov, R. Šajn, S. Pavlov, V. Enimiteva (2012). Trends of atmospheric deposition of trace elements in Macedonia studied by the moss biomonitoring technique, Journal of Environmental Science and Health, Part A, 47(13): 2000-2015.
- L. Barandovski, T. Stafilov, R. Šajn, M. V. Frontasyeva, K. Bačeva, Air pollution study in Macedonia by using moss biomonitoring technique ICP-AES and AAS (2013). Macedonian Journal of Chemistry and Chemical Engineering, 32(2): 265-281.

Romania

Lead institute: Ion V. Popescu, Claudia Stihi, Valahia University of Targoviste, Faculty of Sciences and Arts, Targoviste, Romania. <u>ivpopes@yahoo.com</u>, <u>stihi@valahia.ro</u>
 Collaborating institute(s): Antoaneta Ene, Dunarea de Jos University of Galati, Faculty of Sciences and

Collaborating institute(s): Antoaneta Ene, Dunarea de Jos University of Galati, Faculty of Sciences and Environment, Galati, Romania. <u>aene@ugal.ro</u> Simona Cucu-Man, Alexandru Ioan Cuza University of Iasi, Romania. <u>sman@uaic.ro</u> Radu Todoran, Technical University of Cluj-Napoca, North University Center, Baia Mare, Romania. <u>Todoran_radu@yahoo.com</u> Marina V. Frontasyeva, Otilia Culicov, Joint Institute for Nuclear Research, Dubna, Russian Federation. <u>marina@nf.jinr.ru</u>, <u>culicov@nf.jinr.ru</u>

Background

The first systematic study on atmospheric pollution from heavy metals and other toxic elements based on moss analysis was undertaken as a Romanian–Russian–Norwegian collaboration, in order to assess the general state of heavy metal pollution in Romania in the period 1995–2001 (Lucaciu et al., 2004 and the citations herein). The results on moss samples collected in different regions of Romania in 1990, 1995 and 2000 were unified and reported by Harmens et al. (2008); Romania did not submit data for the 2005/6 European moss survey.

Results and discussion

Nationwide moss survey undertaken in 2010/2011 by four Romanian Universities from Targoviste, Galati, lasi and Baia Mare comprised 330 sampling sites evenly distributed over 75% of the Romanian territory. A total of 34 elements (Na, Mg, Al, Cl, K, Ca, Sc, Ti, V, Cr, Mn, Fe, Co, Ni, Cu*, Zn, As, Br, Rb, Sr, Cd*, Sb, Ba, Cs, La, Ce, Sm, Tb, Hf, Ta, W, Pb*, Th, and U) were determined in the large-scale concentration range — from 10000 mg/kg for Al and K to 0.001 mg/kg for some rare earth elements — by two complementary methods: instrumental epithermal neutron activation analysis (INAA) at the IBR-2 reactor at the Joint Institute for Nuclear Research at Dubna, Russian Federation, and graphite furnace/flame atomic absorption spectrometry (GFAAS /FAAS)^{*} in the Multidisciplinary Research Institute for Science and Technologies from Valahia University of Targoviste, Romania. Principal component (factor) analysis was used to identify the most polluted areas and characterize different pollution sources. Four factors were revealed (**Figure 1**).



Figure 1 Spatial distribution of factor scores based on element concentrations in mosses in Romania in 2010.

Factor 1 is a mixture of light and heavy crust components (Na, Al, Sc, Ti, V, Cr, Fe, Ni, Co, Ba, La, Ce, Sm, Tb, Hf, Ta, W, Th, U). Factor 2 is of anthropogenic origin: Zn (0.82), As (0.73), Sb (0.92), the

main contributors to this factor being mining and industrial sites from north-western parts of the country. **Factor 3** is a mixture of "marine elements" CI (0.79) and Br (0.62); Ca (0.78), K (0.57) and Sr (0.45) may originate from fertilizer components — Sr most probably originates from phosphate fertilizers in agricultural areas and Ca and K from saltpeter (nitre) ones. **Factor 4** comprises Pb (0.63), Cd (0.66), Cu (0.64) which can originate from gasoline or copper mining industry. Although the concentrations of heavy metals in mosses collected in Romania are high compared to other European countries (Harmens et al., 2013), the temporal trends presented in **Figure 2** based on the reported values (Harmens et al., 2008, 2013) for selected metals reveal a decrease for the majority of elements, with the exception of Pb, Cd and Cu.



Figure 2 Temporal trends of metal concentrations in mosses in Romania since 1990.

Conclusion

From the presented results it can be concluded that atmospheric deposition of trace metals is a considerable problem in the northern and western parts of Romania. This study contributes to the national monitoring system of Romania for long range transported elements of air pollutants, and along with epidemiological data it may serve as a baseline for human health risk assessments.

References

- Lucaciu, A., Timofte, L., Culicov, O., Frontasyeva, M.V., Oprea, C., Cucu-Man, S., Mocanu, R., Steinnes, E. (2004). Atmospheric deposition of trace elements in Romania studied by the moss biomonitoring technique, Journal of Atmospheric Chemistry 49: 533–548.
- Harmens, H., Norris, D. & the participants of the moss survey (2008). Spatial and temporal trends in heavy metal accumulation in mosses in Europe (1990-2005). Programme Coordination Centre for the ICP Vegetation, Centre for Ecology and Hydrology, Bangor, UK.
- Harmens, H., Norris, D., Mills, G., and the participants of the moss survey (2013). Heavy metals and nitrogen in mosses: spatial patterns in 2010/2011 and long-term temporal trends in Europe. ICP Vegetation Programme Coordination Centre, Centre for Ecology and Hydrology, Bangor, UK.

Russian Federation

Lead institute: Marina V. Frontasyeva, Joint Institute for Nuclear Research, Dubna, Russian Federation, marina@nf.jinr.ru

Collaborating institutes: Anatoly M. Dunaev, Ivanovo State University of Chemistry and Technology, Ivanovo, amdunaev@ro.ru;

Juliya Koroleva, Immanuel Kant Baltic Federal University, Kaliningrad, <u>vu.koroleff@yandex.ru</u>; Inna V. Vikhrova, Municipal Educational Centre, Tikhhvin, Leningradskaya Region; Nina A. Lebedeva, Liceum # 1, Volgorechensk, Kostromskaya Region.

Background

The sporadic contributions of the Russian Federation to the European moss survey started in the early nineties of the last century in North-West Russia, Sankt-Petersburg, Kaliningrad, Pskov, Novgorod, part of Vologda and part of Karelia (by Goltsova and Fedoretz; Goltsova et al., 1992; Kharin et al., 2001). In 2000, results from Central Russia, Tver, Yaroslavl and Tula Regions were reported (by Ermakova and Frontasyeva; Ermakova et al., 2004). The next moss survey in 2005 was represented by the South Urals, part of the Republic of Udmurtia (by Pankratova and Frontasyeva; Yu et al., 2009) and the region North-East of Moscow Region (by Vergel and Frontasyeva; Vergel et al., 2009). In 2010/2011, moss sampling was carried out in Central Russia, in Ivanovo Region, and on a local scale around the towns of Tikhvin, Lenindradskaya Region, and Volgorechensk, Kostromskaya Region. Apart from maybe Vologda, all sampled areas are affected by industrial activity, such as ferrovanadium plant "Vanadium-Tula" in the city of Tula, ferrochromium smelter "Tikhvin Ferroalloy Plant" in the Tikhvin district known for its bauxite, limestone and dolomite deposits, and many others. A clustering of heavy and machine industries in the Republic of Udmurtia along with oil mining and processing industry severely affect the environment in this area.

Results and discussion

Table 2

The moss survey in 2010/2011 was undertaken by the postgraduate students of Ivanovo State University of Chemistry and Technology and two teachers and their pupils of the secondary schools in Tikhvin and Volgorechensk districts in collaboration with the Joint Institute of Nuclear Research in Dubna, which provided neutron activation analysis (NAA) at the reactor IBR-2 of FLNP JINR. Ivanovo State University of Chemistry and Technology provided atomic absorption spectrometry (AAS) for the samples from their region. The results obtained by AAS from the last moss survey in 2010 in Kaliningrad by Koroleva and colleagues (Koroleva et al., 2012), though not included in the report of the 2010/2011 European moss survey, were presented at the 27th ICP Vegetation Task Force meeting in Paris (**Tables 1 and 2**).

year	Cu	Zn	Mn	Fe	Ni	Pb	Ag	Cr	Cd
1994	5.26	37	-	530	1.85	8.05	-	1.58	0.29
2000	5.03	29	256	192	4.11	13.5	0.13	0.14	0.12
2005	9.59	36	139	220	3.55	4.63	0.04	0.99	0.21
2010	2.84	-	245	168	1.21	1.91	0.03	1.41	0.05

Table 1	Median metal	concentrations	(ma/ka)	in mosses in	the Kaliningrad	Region from	1994-2010
	moulan motal	00110011110110	(1119/119)	1111000000 111	and raining ac	i i togioni nom	1001 2010

	Cu	Ni	Pb	Cr	Cd	Ag	Mn	Fe
mean	4.46	1.20	2.51	2.16	0.109	0.044	346	217
min	1.68	0.36	0.39	0.69	0.006	0.007	73	113
max	19.7	2.54	9.33	12.1	1.12	0.233	960	700

Heavy metal concentrations in *Pleurozium shreberi* in the Kaliningrad Region in 2010 (mg/kg).

The low values (relatively to Norwegian ones) are evidence for the absence of potential sources of heavy metal emissions in Kaliningrad. The highest concentrations were determined in the western part (Sambia peninsula, Baltic Sea coast), elevated levels were observed in the North, North-East and South-West parts, and the lowest ones were in the centre of the Region.

A total of 34 elements (Na, Mg, Al, Cl, K, Ca, Sc, Ti, V, Cr, Mn, Fe, Co, Ni, Cu*, Zn, As, Br, Rb, Sr, Cd*, Sb, Ba, Cs, La, Ce, Sm, Tb, Hf, Ta, W, Pb*, Th, and U) were determined by NAA and AAS (marked with *) in

moss samples collected in Ivanovo region, Tikhvin and Vilgorechensk districts. Some results of moss samples collected from the Ivanovo Region are shown in **Figures 1 and 2**.





Industrial activity is not a characteristic feature for the Ivanovo Region, except for small industrial enterprises in the city of Ivanovo. Therefore, the main contribution in atmospheric deposition is due to transboundary air pollution. The most probable sources of emission of pollutants are situated in the neighbouring regions (Yaroslavl, Kostroma, Vladimir): oil-refinery, metal processing, thermal power plants, and chemical industry. The combined analysis of moss and associated soil allows establishing the main source of anthropogenic impact on the wildlife preserve "Klyazminskii". These are metallurgical plants in Kovrov and Klyazminskii gorodok. Tula and Udmurtia are industrial centres and show the highest concentrations for most of the heavy metals (Figure 2).





An example of the deposition patterns of Fe and Cr, the main contaminants of the town of Tikhvin in Leningradskaya Region, is shown in **Figure 3**.



Figure 3 Distribution maps for iron (left) and chromium (right), the main contaminants of the "Tikhvin Ferroalloy Plant" in the Tikhvin district.

It was shown that the main source of contamination of the environment in the town of Volgorechensk of Kostromskaya Region is the thermal power plant allocated in the suburbs of the town. Multivariate statistical analysis (PCA) was applied to the analytical results obtained for all samples areas to reveal and characterize the pollution sources.

Conclusions

It follows from the results obtained in the moss survey 2010/2011 in the sampled areas under different anthropogenic loadings of pollutants that contamination with heavy metals and other toxic elements is rather local but it may lead to the environmental stress. Attempts were made to use the results in environmental and human health risk assessments.

References

- N.I. Goltsova, T.V. Vasina (1992). Heavy metal content in fores moss Pleurozium schreberi in Leningradskaya Region. Environmental Chemistry 2:.27.
- V. N. Kharin, N.G. Fedorets, G V. Shiltsova, V.V. Dyakonov, E.N. Spektor (2001). Geographic trends in the accumulation of heavy metals in mosses and forest litters in Karelia. Russian Journal of Ecology 32: 138–141. Translated from Ekologiya 2: 155–158.
- E.V. Ermakova, M.V. Frontasyeva, S.S. Pavlov, E.A. Povtoreyko, E. Steinnes, Ye.N. Cheremisina (2004). Air pollution studies in Central Russia (Tver and Yaroslavl Regions) using the moss biomonitoring technique and neutron activation analysis. Journal of Atmospheric Chemistry 49: 549-56.
 K.N. Vergel, M.V. Frontasyeva, I.Z. Kamanina, S.S. Pavlov (2009). Biomonitoring of atmospheric heavy metal deposition in the
- K.N. Vergel, M.V. Frontasyeva, I.Z. Kamanina, S.S. Pavlov (2009). Biomonitoring of atmospheric heavy metal deposition in the north-east of Moscow Region by means of moss biomonitoring technique. Ecology of Urbinized Terrotories (RF) 3: 88-95 (in Russian).
- Yu. S. Pankratova, M.V. Frontasyeva, S.S. Pavlov, N.I. Zelnitchenko (2009). Atmospheric pollution in the Republic of Udmutria – estimations based on moss analysis. Problems of Regional Ecology 1: 57-63, c. 99-107.
- Koroleva Y.V., Barinova G.M., Krasnov E. V. (2012). Indicative Modeling and Spatial Evaluation of Air Pollution Risk: Risk Models and Application, Collected Paper, Co-Data - Germany, Berlin, p. 23-34.

Serbia

Lead institute: Marko Sabovljević; Faculty of Biology, University of Belgrade, Serbia. <u>marko@bio.bg.ac.rs</u> Collaborating institutes:

Mitja Skudnik; Slovenian Forestry Institute, Ljubljana, Slovenia. <u>mitja.skudnik@gozdis.si</u> Zvonka Jeran;Institute Jožef Stefan, Ljubljana, Slovenia. <u>zvonka.jeran@ijs.si</u> Franc Batič; Biotechnical Faculty, University of Ljubljana, Slovenia. franc.batic@bf.uni-lj.si

Background

Serbia was involved in the European moss survey for the first time in 2005, when the atmospheric deposition of trace elements was studied mainly in the northern part of the country. In 2012, the second moss survey was performed in Serbia (18 different trace elements, nitrogen and sulphur), but for the first time on the national scale (excl. Kosovo). Mosses were sampled in a one-month period (mid May - mid June) for 42 locations that were regularly distributed throughout the country inside the systematic sampling grid UTM 50 km x 50 km. At each of the plots 5 subsamples of the moss *Hypnum cupressiforme* Hedw. were collected. Field sampling was done according to the guidelines of the UNECE ICP Vegetation programme. With the aim to achieve rather representative values of deposit in mosses per plot, one composite sample was pooled from five subsamples at each location. Before analysis, mosses were dried at room temperature and cleaned from obvious contamination particles (soil particles, litter etc.). Only green and yellow-green parts of the mosses were taken for further analysis. Moss materials was lyophilized for 24 hours and homogenized in a ball mill. For each of the most important pollution emission sources within the county are industry, smelters, mining and traffic.

Results

Element concentrations in mosses are shown in **Tables 1 and 2**. Results indicate higher contents of arsenic and chromium in moss tissues in Serbia. In the eastern part, the region of the Bor smelter is especially loaded with cadmium and copper. Nickel is elevated in the western part of the country, probably due to industry. The mean values of trace elements in the moss tissues in Serbia are somewhat higher than those reported for some western European countries, but more or less in accordance with neighbouring areas. Nitrogen deposition is especially high in the areas around bigger cities (Belgrade, Niš), along the Sava river and in the northern part of the country. Sulphur deposition is high in northern Serbia and areas around bigger cities, especially Belgrade and Niš.

[µg/g]	As	Cd	Cr	Cu	Fe	Ni	Pb	Sb	V	Zn
Min	0.22	0.10	1.31	3.99	481.99	2.16	2.56	0.05	1.43	18.42
Max	6.50	0.99	23.92	78.70	9817.46	37.58	42.26	1.18	20.58	87.00
mean	1.67	0.27	6.26	11.92	2351.55	6.93	10.54	0.19	6.10	41.58
median	1.28	0.22	5.24	9.31	2004.39	4.65	7.76	0.14	5.42	34.51

Descriptive statistics of heavy metal content in mosses in Serbia in 2012 (N=42).

Table 2

Table 1

Descriptive statistics of N and S contents in mosses in Serbia in 2012 (N=42).

[mg/g]	Ν	S
min	9.04	0.84
max	28.73	2.63
mean	16.65	1.45
median	14.25	1.34

The spatial patterns of the lead (Pb) and nitrogen (N) content in mosses in Serbia in 2012 is shown in **Figure 1**.





Spatial patterns of Pb (left) and N (right) content in the moss *Hypnum cupressiforme* in Serbia in 2012 per 50 km x 50 km grid.

Discussion, conclusions, recommendations

The spatial variability of trace elements, nitrogen and sulphur, accumulated in mosses correlates well with the spatial variability in the level of the industrialization, traffic intensity and other anthropogenic activities. Because the analysis was done for the first time on the national scale, we cannot report on temporal trends. The abundance of the moss *Hypnum cupressiforme* is not equal in the whole territory of Serbia. In some areas (especially in northern parts of the country) it was difficult to collect five sub-samples at suitable sites. The monitoring with some other moss species should be regarded as an option. Microhabitat characteristics could be the reason for slightly differences in deposit values, due to the differences to precipitate exposures and further studies are needed. Individual and regional efforts, so far, do not allow for extensive analyses, such as long-term spatial deposition changes, spatial pattern changes or temporal national or transboundary patterns. Both research and/or screening as well as permanent monitoring are needed. National monitoring systems of atmospheric deposition including moss survey should be established long term as a *National air quality control programme*.

The authorities should establish a national network for atmospheric pollution biomonitoring, as well as national permanent sampling plots. Moss monitoring with at least the current sampling density, expanded to include more atmospheric pollutants and new technologies should be established. The nationwide moss survey should be conducted frequently, providing a detailed record of spatial patterns and temporal trends of atmospheric deposition over Serbia. This is an important and valuable supplement to the national monitoring of trace metals in precipitation, which is limited to a small number of sites.

Slovenia

Lead institutes: Franc Batič, Boris Turk, University of Ljubljana, Biotechnical Faculty, Agronomy Department., Jamnikarjeva 101, 1000 Ljubljana, Slovenia; <u>franc.batic@bf.uni-lj.si</u>, <u>boris.turk@bf.uni-lj.si</u> (ozone research).

Żvonka Jeran, Darja Mazej, Jožef Stefan Institute, Department of Environmental Sciences, Ljubljana, Slovenia; <u>zvonka.jeran@ijs.si</u>, <u>darja.mazej@ijs.si</u> (moss survey).

Collaborating institutes: Nataša Kopušar Institute for Ecological Research Velenje, Koroška cesta 58, 3320 Velenje; <u>natasa.kopušar@erico.si</u> (ozone research).

Mitja Skudnik, Daniel Žlindra, Slovenian Forestry Institute, Ljubljana, Slovenia; <u>mitja.skudnik@gozdis.si</u>, <u>Daniel.zlindra@gozdis.si</u> (moss survey).

Background

Slovenia has been involved in the European moss survey since 1995 and in ozone research since 1996. Mosses have been collected every five years between the middle of June and the end of July on a regular grid. The following elements were determined: As, Cd, Cr, Co, Cu, Fe, Hg, Mo, Ni, Pb, Sb, Se, Sr, V, Ti and Zn. Since 2001, nitrogen and sulphur were included, and in 2010 the first results for polycyclic aromatic hydrocarbons (PAHs) in mosses were obtained from a selected number of plots. Regarding ozone, the main objective is to monitor effects of tropospheric ozone on crops and semi-natural vegetation in agreement with the Slovenian National Environmental Program. Three permanent experimental sites were established by the group in Ljubljana representing air pollution and land use types in Slovenia: Ljubljana (urban site with moderate level of ozone), Iskrba (rural site, with elevated level of ozone) and Rakican (EMEP station, rural site important for crop production, affected by high traffic) and several sites around the thermal power plant Šoštanj established by the group in Velenje. In the years 1996-2007, white clover (*Trifolium repens* 'Regal') was used as indicator species, then brown knapweed (*Centaurea jacea*) and bush bean (*Phaseolus vulgaris*) were introduced as ozone indicators in 2008-2010 following the ICP Vegetation protocols.

The emissions of ozone precursors in Slovenia are showing 41% decline from 1990 to 2011. NO_x emissions declined 25%, CO emissions 56%, non-methane VOCs 51% and methane emissions 7%. The main reason for the decline is introduction of new emission standards for motor vehicles. Additional reasons for the decline are new legislation on storing and transport of fuels, technological improvements in thermal power plants, installation of exhaust cleaning devices and improvement of incineration processes in industry. Tropospheric ozone concentration shows distinct fluctuation during the year, and among the years, depending on sunlight. They are the highest in the western parts of Slovenia, due to the transboundary transport of precursors from northern Italy. Measured ozone concentrations are mostly above target values, all over the country. No significant trend can be observed in long-term measurements of ozone concentrations.

Results

Ozone. Tropospheric ozone concentrations, expressed as AOT40 values, are showing fluctuation with time, mostly depending on weather conditions and thus not following the general ozone precursors decline, discussed above (**Figure 1**). Due to transboundary inflow of precursors and higher elevation, significantly higher concentration are observed at the lskrba site.





AOT40 calculated at three sites in Slovenia between 1998 and 2007.
Observed ozone-induced leaf injuries of white clover (**Figure 2**) showed some correlation with measured AOT40 values of ozone, but are confounded by other environmental influences (drought, temperature, pests) and the subjective nature of assessment (different assessors with time and location). White clover proved not to be a reliable indicator of ozone impacts on plants, at least at such small scale, bush bean seems to be slightly more reliable (data not shown).



Figure 2

Ozone-induced leaf injury on white clover for an ozone sensitive and resistant clone.





Concentrations of cadmium and nitrogen in *H. cupressiforme* collected at 102 locations in Slovenia in 2010 (left) and concentrations of cadmium and nitrogen at 32 plots which were the same in the last three surveys, i.e. 2001, 2006 and 2010 (right).

Heavy metals and nitrogen. The overall spatial pattern (**Figure 3**) of levels of anthropogenically derived elements especially Pb, Cd, Mo, Zn, Sb and N in Slovenia can be explained on the one hand by the population density and the locations of the main pollution sources (steel factories at Jesenice and Ravne, coal burning thermal power plants at Šoštanj and Trbovlje, a former Pb-smelter) and on the other hand by transboundary transport of air pollutants from neighbouring countries, mainly as a consequence of meteorological conditions (wind, precipitation). The northern and north-western parts of Slovenia are the regions receiving the highest yearly amounts of precipitation, while the precipitation pattern gradually decreases toward the east. The heavy transit traffic that crosses

Slovenia along the diagonal from the port of Koper or Gorica in the south-west to the Hungarian or Austrian border in the north-east might be an additional source of traffic-derived metals such as Sb, as well as for N. Higher values of N in moss in the north-eastern region can additionally point to the agricultural activity that dominates this area.

Similar to the majority of European countries, a decrease of heavy metal concentrations in mosses was observed in Slovenia since 1995 (**Figure 4**), which was most probably due to economic factors because some of the heavy polluters have shut down or modernized their technology to reduce emissions. Another reason for the relatively big decline between 2005 and 2010 could be a change in sampling strategy. Influence of canopy drip was avoided in 2010. Comparative analyses had shown that significantly higher concentrations were found in mosses under the canopy in comparison to those at least 1 m away; on average about 20% for Cu, As and Hg, to up to 30- 50% for Cu, Hg, Ni, V, Pb and N 3 m away from the nearest tree.





Discussion, conclusions, recommendations

Biomonitoring of tropospheric ozone effects in Slovenia, carried out within the activity of ICP-Vegetation, has proven that there is a high potential for ozone formation in the country but the observed damage on bioindicator plants used was much smaller than expected, based on AOT values. The reason for this discrepancy might be over-sensibility of used indicator plants to drought stress, especially white clover, what was proven by some diurnal measurements of photosynthesis and transpiration at the site in Ljubljana. We got similar results with bush bean when the plants were covered with a hail protection net. The net reduced sunlight intensity and typical ozone damages were observed in both biotypes of plants. High potential risk of ozone damage on plants was also proven by ozone biomonitoring with tobacco (*Nicotiana tabacum* 'Bel-W3') and the high frequency of ozone damage on forest vegetation and ornamental plants. Biomonitoring of ozone impacts on vegetation as developed within activities of Working Group on Effects of the LRTAP Convention should be integrated in the future in European environmental policy and common agro-environmental programmes to ensure continuity and enhance chances of funding.

Although there were slight modifications in moss sampling over the 15 year period, the spatial pattern of elements and their temporal trends reflect the overall pollution in the country, as well as the natural background. The application of factor analysis to the moss data resulted in similar elemental associations in a particular factor and its spatial distribution in all three surveys, including a lichen survey (Jeran et al., 2007). Moss monitoring should be used in the future as a supplement to the official national air monitoring programmes performed by the Slovenian Environment Agency, as it could provide additional data on air pollution (elements, nitrogen, organic pollutants and radionuclides) at a higher number of sites than covered by their current programmes.

References

Jeran, Z., Mrak, T., Jaćimović, R., Batič, F., Kastelec, D., Mavsar, R., Simončič, P. (2007). Epiphytic lichens as biomonitors of atmospheric pollution in Slovenian forests. Environmental pollution 146: 324-331.



Air Pollution: Deposition to and impacts on vegetation in (South-)East Europe, Caucasus, Central Asia (EECCA/SEE) and South-East Asia

This report is a synthesis of current knowledge on the deposition of air pollutants to terrestrial ecosystems and their impacts on vegetation in countries in Eastern Europe, Caucasus and Central Asia (EECCA), South-East Europe (SEE) and South-East Asia (Malé Declaration countries). Pollutants included are primarily ozone, nitrogen and heavy metals but also persistent organic pollutants (POPs) for EECCA/SEE countries and black carbon (as a component of aerosol) for South-East Asia. One of the current priorities of the Convention on Long-range Transboundairy Air Pollution (LRTAP) is the ratification of its Protocols by EECCA countries, with the aim to reduce air pollution in those countries and in neighbouring countries from long-range transport. In recent decades, the decline in air pollution in EECCA and SEE countries has generally been lower than the reductions achieved in Northern, Western and Central Europe. Furthermore, the Convention is keen to stimulate outreach activities to regions outside the UNECE area. Hence the focus of this review on the above-mentioned countries.

For further information or copies contact: Harry Harmens Centre for Ecology and Hydrology Environment Centre Wales Deiniol Road Bangor Gwynedd LL57 2UW United Kingdom Tel.: +44 (0) 1248 374500 Fax: +44 (0) 1248 362133 Email: hh@ceh.ac.uk

ICP Vegetation web site: http://icpvegetation.ceh.ac.uk



ISBN: 978-1-906698-48-5

